

**NOTES SUR LES TECHNOLOGIES DE LA SANTÉ EN  
ÉMERGENCE**

# Anticorps monoclonaux pour prévenir les migraines

## Résumé

- La migraine est un trouble neurologique commun et chronique. La toxine botulique a reçu l'approbation réglementaire pour la prévention de la migraine chronique. Ce médicament nécessite de multiples injections à des points précis d'injection à la tête et au cou. D'autres médicaments sont utilisés dans la prévention de la migraine, mais l'adhésion des patients et l'efficacité demeurent problématiques.
- Les anticorps monoclonaux antagonistes des récepteurs du peptide lié au gène de la calcitonine (anti-CGRP) appartiennent à une nouvelle classe de médicaments qui ciblent le CGRP — un puissant vasodilatateur — qui joue un rôle dans la douleur et les migraines.
- Quatre anticorps monoclonaux anti-CGRP sont en cours de développement pour être utilisés dans la prévention de migraines épisodiques et chroniques : l'éptinezumab, l'érenumab, le frémanézumab, et le galcanézumab. À ce jour, ils n'ont été approuvés dans aucun pays.
- Des résultats d'essais de phase II et de phase III sont diffusés pour les quatre médicaments. Les données probantes disponibles à ce jour présentent une diminution statistiquement significative de la fréquence des migraines. Les individus souffrant de migraines épisodiques ont eu un à deux épisodes (jours) de migraine de moins par mois. Les patients souffrant de migraines chroniques ont eu de deux à deux épisodes (jours) et demi de migraine de moins par mois. Aucune préoccupation concernant l'innocuité de ces médicaments n'a été relevée.
- Bien que le coût de ces médicaments n'ait pas été défini, ils auront potentiellement un important impact budgétaire étant donné le coût élevé des produits biologiques et la prévalence des migraines épisodiques et chroniques.

## Background

A migraine is characterized as a headache, with at least two of the following pain attributes: moderate or severe, throbbing, localized to one area, and avoidance of routine physical activity because of the pain.<sup>1,2</sup> The headache must be accompanied by at least one of the following symptoms: nausea or vomiting, or photophobia (light sensitivity) and phonophobia (sensitivity to noise).<sup>1</sup> Up to 20% of patients with migraines may experience a visual or auditory disturbance (aura); a migraine attack can last from four hours to 72 hours.<sup>3,4</sup>

Migraines are classified, based on the frequency of attacks, into episodic migraine (headaches that occur less than 15 days per month) and chronic migraine (headaches for 15 days or more per month).<sup>5,6</sup> In Western countries, episodic migraines affect approximately 12% of the adult population, while chronic migraines affect approximately 1% of the adult population.<sup>2,4,5</sup> Chronic migraines affect three times more women than men and are associated with an increased societal burden, an increased number of comorbidities, and a negative impact on quality of life.<sup>2,4,5</sup>

Research is in progress to develop drugs specifically for the prevention of episodic and chronic migraines. It is reported that migraines are the third most-prevalent and the seventh most-disabling condition worldwide.<sup>7</sup> Therefore, these medications will potentially be important in improving the quality of life and reducing the disease burden of patients who suffer from chronic or episodic migraines.

## The Technology

Calcitonin gene-related peptide (CGRP) is a 37 amino acid neuropeptide present in the peripheral and central nervous systems.<sup>6,8</sup> A release of CGRP has been shown to induce migraine attacks.<sup>9</sup> Anti-CGRP monoclonal antibodies act to weaken the migraine signalling pathway.<sup>9</sup>

Previous studies conducted on small molecules targeting the CGRP receptors demonstrated efficacy in treating and preventing migraines. However, these trials were stopped prematurely because of serious concerns of liver toxicity and other reasons (Appendix 1).<sup>6,8</sup>

Monoclonal antibodies (mAbs) targeting the CGRP signalling pathway are currently being developed as preventive therapy for episodic and chronic migraines.<sup>3,6,8</sup> Clinical trials are in progress for three mAbs targeting CGRP and one targeting the CGRP receptor.<sup>3,6,8</sup> Anti-CGRP mAbs remove excess CGRP molecules, while anti-CGRP receptor mAbs block signalling at the receptor.<sup>6</sup> This prevents the activation of CGRP signalling pathways, which may decrease headache frequency over time.<sup>3,6</sup>

The three mAbs targeting CGRP are eptinezumab (ALD403; Alder Biopharmaceuticals), fremanezumab (TEV-48215 or LBR-101; Teva Canada Innovation), and galcanezumab (LY2951742; Eli Lilly Canada Inc.), while erenumab (AMG 334; Amgen Canada) is targeting the CGRP receptor.<sup>3,6,8</sup>

## Regulatory Status

These drugs have not yet been approved to be marketed in any country.

*Eptinezumab*: Alder Biopharmaceuticals will be making a Biologics License Application (BLA) to the FDA for eptinezumab in the second half of 2018.<sup>10</sup>

*Erenumab*: Erenumab is currently under review by Health Canada.<sup>11</sup> On June 21, 2017, the European Medicines Agency accepted a marketing authorization application from Novartis for erenumab.<sup>12</sup> The FDA accepted a BLA for erenumab on July 20, 2017.<sup>13</sup> Of note, Amgen is transitioning the Canadian commercialization rights of erenumab to Novartis (Geoff Sprang, Executive Director, Value & Access, Amgen Canada, Mississauga, ON: personal communication, 2017 Sep 11).

*Fremanezumab*: Teva announced the submission of a BLA to the FDA for fremanezumab on October 17, 2017.<sup>14</sup> Submissions to other regulatory agencies will follow at a later date. (Christine Poulin, General Manager, Teva Canada Innovation, Montreal, QC: personal communication, 2017 Sep 13).

*Galcanezumab*: Eli Lilly has submitted a BLA for galcanezumab to the FDA;<sup>15</sup> submissions to other regulatory agencies will occur at a later date (Doron Sagman, Senior Medical Director, Eli Lilly Canada Inc., Toronto, ON: personal communication, 2017 Sep 13).

## Cost and Administration

The costs for erenumab, fremanezumab, and galcanezumab are currently unavailable (Doron Sagman : personal communication, 2017 Sep 13; Christine Poulin: personal communication, 2017 Sep 13; Geoff Sprang: personal communication, 2017 Sep 11). We have no information on eptinezumab.

All four drugs are administered parenterally; dosage information obtained from the clinical trials is available in Table 1, although these may change based on Health Canada's regulatory reviews.

**Table 1: Administration of Anti-CGRP Monoclonal Antibodies**<sup>10,16-23</sup>

Drug	Dose(s)	Route of Administration	Dosing Interval
Eptinezumab	100 mg or 300 mg	intravenous infusion	every 3 months
Erenumab	70 mg or 140 mg	subcutaneous injection	every month
Fremanezumab	225 mg	subcutaneous injection	every month
	675 mg	subcutaneous injection	every 3 months
Galcanezumab	120 mg or 240 mg	subcutaneous injection	every month

anti-CGRP = anti-calcitonin gene-related peptide.

<sup>a</sup> 675 mg initial dose followed by 225 mg monthly dose in chronic migraines.

## Target Population

The intended use of these anti-CGRP mAbs is for migraine prevention in patients suffering from episodic or chronic migraines.<sup>1,24</sup> Patients who have failed other preventive migraine therapies may be a target population for the use of anti-CGRP therapy.<sup>8,25</sup>

## Current Practice

Existing therapies for migraine management focus on both the acute and the preventive management of symptoms.<sup>1,24</sup> 5HT<sub>1B/1D</sub> agonists, known as triptans, are the mainstay for the treatment of acute migraine episodes and are available in a number of formulations.<sup>1,24</sup> Other pharmacologic options for treating acute migraines include analgesics (acetaminophen) and nonsteroidal anti-inflammatory drugs (ibuprofen, naproxen, and ASA).<sup>24</sup>

Patients who suffer from recurrent attacks may benefit from prophylactic therapy.<sup>26</sup> Botulinum toxin has received regulatory approval for the prevention of chronic migraine headaches. This medication requires multiple injections into specific head and neck sites.<sup>1,6,24</sup> Topiramate is used for the prophylaxis of migraine headaches in adults experiencing four or more migraine attacks per month.<sup>27</sup> Other drugs used for migraine prevention are anti-epileptics (divalproex sodium, lamotrigine, gabapentin), antidepressants (amitriptyline, nortriptyline, fluoxetine, venlafaxine, duloxetine), beta blockers (metoprolol, propranolol), calcium channel blockers (verapamil, flunarizine), and magnesium. These drugs have not been approved for use in migraine prevention and are associated with low adherence because of side effects or efficacy.<sup>1,5,8,24,26,28</sup>

# Methods

These bulletins summarize information available about a new or emerging technology. Industry is invited to review the information, and drafts are reviewed by at least one clinical expert.

A peer-reviewed literature search was conducted using the following bibliographic databases: MEDLINE, PubMed, Embase, and the Cochrane Library. Grey literature was identified by searching relevant sections of the *Grey Matters* checklist (<https://www.cadth.ca/grey-matters>). No methodological filters were applied. The search was not limited by language or publication date. Regular alerts updated the search until project completion; only citations retrieved before 18 December 2017 were incorporated into the analysis. Conference abstracts were included from the search results.

## Summary of the Evidence

### Episodic Migraine

A total of 10 randomized, double-blind, placebo-controlled trials have been conducted in adult patients: three of these trials were phase II trials, one was a phase IIb trial, and six were phase III trials. All trials involved multiple testing centres and all were industry-sponsored (Table 2). The sample sizes ranged from 174 patients to 955 patients. The patient population included adult patients experiencing episodic migraines for 12 months or longer before screening. Patients were randomly assigned to receive either the treatment drug or placebo. The primary outcome measured for these drugs was the change in the frequency of migraines. The trial durations were 24 weeks for the PROMISE-1, STRIVE, EVOLVE-1, and EVOLVE-2 studies; and 12 weeks for the other trials.

Author (Year) Name of Study, Country, Funding	Study Phase, Study Duration, Sample Size	Population	Intervention(s), Comparator	Primary Outcome
<b>Episodic Migraine</b>				
<b>Eptinezumab</b>				
Alder Biopharmaceuticals Press Release (2017) <sup>10</sup>  ClinicalTrials.gov (NCT02559895) <sup>29</sup>  Keller (2017) <sup>30</sup>  PROMISE-1 Multi- centre, US  Alder Biopharmaceuticals	Phase III  24 weeks  N = 888	Patients 18- to 65- years-old with episodic migraines	Eptinezumab <sup>a</sup> 100 mg IV every 3 months (n = 222)  Eptinezumab 300 mg IV every 3 months (n = 222)  Placebo (n = 222)	Change from baseline over weeks 1 to 12 in monthly migraine days
Dodick et al. (2014) <sup>7</sup>  Multi-centre, US  Alder Biopharmaceuticals	Phase II  12 weeks  N = 174	Patients 18- to 55-year- old with 5 to 14 migraine days per month	Eptinezumab 1,000 mg IV one dose (n = 86)  Matching placebo (n = 88)	Change from baseline in weeks 5 to 8 in the frequency of migraine days
<b>Erenumab</b>				

Dodick et al. (2017) <sup>16</sup> ClinicalTrials.gov (NCT02483585) <sup>31</sup> Brauser (2017) <sup>32</sup> ARISE North America and Europe Amgen	Phase III 12 weeks N = 577	Patients 18- to 65-years-old with 4 to 14 migraine days per month	Erenumab 70 mg SC monthly (n = 289) Placebo (n = 291)	Change from baseline to weeks 9 to 12 in monthly migraine days
Goadsby et al. (2017) <sup>17,18,33</sup> ClinicalTrials.gov (NCT02456740) <sup>34</sup> Brauser (2017) <sup>32</sup> STRIVE North America and Europe Amgen	Phase III 24 weeks N = 955	Patients 18- to 65-years-old with 4 to 14 migraine days per month	Erenumab 70 mg SC monthly (n = 317) Erenumab 140 mg SC monthly (n = 319) Placebo (n = 319)	Change from baseline in monthly migraine days over weeks 13 to 24
Sun et al. (2016) <sup>35</sup> North America and Europe Amgen	Phase II 12 weeks N = 483	Patients 18- to 60-years-old with 4 to 14 migraine days per month	Erenumab <sup>b</sup> 70 mg SC monthly (n = 107) Matching placebo (n = 160)	Change from baseline in monthly migraine days between weeks 9 to 12
<b>Fremanezumab</b>				
Teva Press Release (2017) <sup>20,21</sup> Keller (2017) <sup>30</sup> HALO EM Multi-centre (global) Teva Pharmaceutical Industries	Phase III 12 weeks N = 875	Patients 18- to 70-years-old with 6 to 14 days per month	Fremanezumab 225 mg SC for 3 months (monthly dose regimen) (n = 290) Fremanezumab 675 mg at initiation, followed by placebo for 2 months (quarterly dose regimen) (n = 291) Matching placebo, 3 monthly doses (n = 294)	Change from baseline in the monthly migraine days during the 12-week period
Bigal et al. (2015) <sup>22</sup> Multi-centre, US Teva Pharmaceutical Industries	Phase IIb 12 weeks N = 297	Patients 18- to 65-years-old with 8 to 14 migraine days per month	Fremanezumab 225 mg SC monthly (n = 96) Fremanezumab 675 mg SC monthly (n = 97) Matching placebo (n = 104)	Change from baseline in the number of migraine days during weeks 9 to 12
<b>Galcanezumab</b>				

Eli Lilly Press Release (2017) <sup>19</sup> ClinicalTrials.gov (NCT02614183) <sup>36</sup> Keller (2017) <sup>30</sup> EVOLVE-1 US, Canada Eli Lilly	Phase III 24 weeks N = 825	Patients 18- to 65-years-old with migraines 8 to 14 days per month	Galcanezumab 120 mg SC monthly (n = NR) Galcanezumab 240 mg SC monthly (n = NR) Matching placebo (n = NR)	Change from baseline in monthly migraine days over 24 weeks
Eli Lilly Press Release (2017) <sup>19</sup> ClinicalTrials.gov (NCT02614196) <sup>37</sup> EVOLVE-2 Multi-centre (global) Eli Lilly	Phase III 24 weeks N = 825	Patients 18- to 65-years-old with migraines 8 to 14 days per month	Galcanezumab 120 mg SC monthly (n=NR) Galcanezumab 240 mg SC monthly (n=NR) Matching placebo (n=NR)	Change from baseline in monthly migraine days over 24 weeks
Dodick et al. (2014) <sup>38</sup> Multi-centre, US Arteaus Therapeutics	Phase II 12 weeks N = 218	Patients 18- to 65-years-old with 4 to 14 migraine days per month	Galcanezumab 150 mg SC injection every 2 weeks (n = 108) Placebo (n = 110)	Change from baseline to weeks 9 to 12 in monthly migraine days

EM = episodic migraine; IV = intravenous; NR = not reported; SC = subcutaneous.

<sup>a</sup> A dose of 30 mg (n = 222) was also tested in the trial.

<sup>b</sup> Doses of 7 mg (n = 108) and 21 mg (n = 108) were also tested in the trial.

## Chronic Migraine

A total of five randomized, double-blind, placebo-controlled trials have been conducted in adult patients: two were phase II trials, one was a phase IIb trial, and two were phase III trials. All trials involved multiple testing centres and all were industry-sponsored (Table 3). The sample sizes ranged from 264 patients to 1,130 patients. The patient population included adult patients with a history of chronic migraine. Patients were randomized to receive either the treatment drug or placebo. The drugs were tested over a 12-week period. The primary outcome included the change in frequency of migraines or headaches.

**Table 3: Characteristics of Randomized, Double-Blind, Placebo-Controlled Trials Chronic Migraine**

Author (Year) Name of Study, Country, Funding	Study Phase, Study Duration, Sample Size	Population	Intervention(s), Comparator	Primary Outcome
<b>Chronic Migraine</b>				
<b>Eptinezumab</b>				

Smith et al. (2017) <sup>39</sup> ClinicalTrials.gov (NCT02275117) <sup>40</sup> US, Australia, New Zealand, Georgia Alder Biopharmaceuticals	Phase II 12 weeks N = 665	Patients 18- to 55-years-old with 15 to 28 headache days per month (at least 8 migraine days)	Eptinezumab <sup>a</sup> 100 mg IV one dose (n = NR) Eptinezumab 300 mg IV once (n = NR) Matching placebo (n = NR)	% patients achieving 75% reduction from baseline in migraine days over weeks 1 to 12
<b>Erenumab</b>				
Tepper et al. (2017) <sup>41</sup> North America and Europe Amgen	Phase II 12 weeks N = 667	Patients 18- to 65-years-old with 15 or more headache days per month (at least 8 migraine days)	Erenumab 70 mg SC monthly (n = 191) Erenumab 140 mg SC monthly (n = 190) Matching placebo (n = 286)	Change from baseline to weeks 9 to 12 in monthly migraine days
<b>Fremanezumab</b>				
Silberstein et al. (2017) <sup>42</sup> Teva Press Release (2017) <sup>20,21</sup> ClinicalTrials.gov (NCT02621931) <sup>43</sup> HALO-CM Multi-centre Teva Pharmaceutical Industries	Phase III 12 weeks N = 1,130	Patients 18- to 70-years-old with a history of chronic migraine	Fremanezumab 675 mg SC followed by monthly 225 mg for two months (monthly dosing regimen) (n = 379) Fremanezumab 675 mg SC at initiation followed by placebo for two months (quarterly dosing regimen) (n = 376) Matching placebo (n = 375)	Change from baseline in monthly average number of headache days of at least moderate severity over 3 months after the first dose of study drug
Bigal et al. (2015) <sup>23</sup> Multi-centre, US Teva Pharmaceutical Industries	Phase IIb 12 weeks N = 264	Patients 18- to 65-years-old with chronic migraine	Fremanezumab 675 mg first dose, then 225 mg <sup>b</sup> SC monthly (n = 88) Fremanezumab 900 mg SC monthly (n = 87) Matching placebo (n = 89)	Change from baseline in headache-hours during weeks 9 to 12
<b>Galcanezumab</b>				
Eli Lilly Press Release (2017) <sup>19</sup> ClinicalTrials.gov (NCT02614261) <sup>44</sup> Keller (2017) <sup>30</sup> REGAIN Multi-centre Eli Lilly	Phase III 12 weeks N = 825	Patients 18- to 65-years-old with at least 15 migraine headaches days per month	Galcanezumab 120 mg SC monthly <sup>c</sup> (n = 278) Galcanezumab 240 mg SC once monthly <sup>c</sup> (n = 277) Matching placebo (n = 558)	Change from baseline in monthly migraine days over 12 weeks

IV = intravenous; NR = not reported; SC = subcutaneous.

<sup>a</sup> Doses of 10 mg and 30 mg were also tested in the trial.

<sup>b</sup> Patients in this trial arm received an initial dose of 675 mg, followed by monthly doses of 225 mg.

<sup>c</sup> Participants received an initial dose of 240 mg.

# Results

## Efficacy

The results of the primary outcomes of the phase II/Ib and phase III trials are depicted in Tables 4 and 5. Of note, except for the STRIVE and HALO-CM studies, the results of the phase III trials have not yet been fully published; the data were obtained from conference abstracts and press releases.

Overall, there was a statistically significant decrease in the frequency of migraines compared with placebo. Episodic migraine sufferers experienced one to two fewer migraine days each month. Patients with chronic migraines experienced two to two-and-one-half fewer migraine days each month.

**Episodic Migraine:** The primary outcome measured in the trials for episodic migraine was the change from baseline in monthly migraine days.

*Eptinezumab:* In the PROMISE-1 study, the mean differences in monthly migraine days for eptinezumab 100 mg and eptinezumab 300 mg were 0.7 days ( $P = 0.0179$ ) and 1.1 days ( $P = 0.0001$ ) compared with placebo, respectively.

In a phase II trial, a single dose of eptinezumab 1,000 mg resulted in a mean difference of 1 day (95% confidence interval [CI], 2.1 to 0.1,  $P = 0.0306$ ), with migraine in weeks 5 to weeks 8 compared with placebo.

*Erenumab:* In ARISE and in STRIVE, the mean differences in monthly migraine days for erenumab 70 mg were 1.1 days ( $P < 0.001$ ) and 1.4 days ( $P < 0.001$ ) compared with placebo, respectively. At the 140 mg dose, the mean difference was 1.9 days ( $P < 0.001$ ) in the STRIVE study.

In a phase II trial, the mean difference in monthly migraine days was 1.1 days (95% CI, 2.1 to 0.2,  $P = 0.021$ ) when comparing erenumab 70 mg with placebo.

*Fremanezumab:* The mean difference in monthly migraine days for fremanezumab 225 mg compared with placebo was 1.5 days ( $P < 0.0001$ ) in the HALO-EM study. The mean difference in the number of migraine days during weeks 9 to 12 was 2.81 days (95% CI, 4.07 to 1.55,  $P < 0.0001$ ) in the phase IIb trial.

Comparing a single dose of fremanezumab 675 mg with a placebo quarterly dose regimen, the mean difference in monthly migraine days was 1.2 days ( $P < 0.0001$ ) in HALO-EM. The mean difference in the number of migraine days during weeks 9 to 12 was 2.64 days (95% CI, 3.90 to 1.38,  $P < 0.0001$ ) for fremanezumab 675 mg compared with placebo in the phase IIb trial.

*Galcanezumab:* In EVOLVE-1 and in EVOLVE-2, the mean differences in monthly migraine days for galcanezumab 120 mg were 1.9 days ( $P < 0.001$ ) and 2.0 days ( $P < 0.001$ ) compared with placebo, respectively. At the 240 mg dose, the mean differences were 1.8 days ( $P < 0.001$ ) and 1.9 days ( $P < 0.001$ ) compared with placebo, respectively.

In a phase II trial, the mean difference in monthly migraine days was 1.2 days (95% CI, 1.9 to 0.6,  $P = 0.003$ ) when comparing galcanezumab 150 mg with placebo.

**Table 4: Change in Frequency of Migraines<sup>a</sup> Episodic Migraine**

Eptinezumab	Phase III Trial (PROMISE-1) <sup>10,30</sup>		
<b>Change from baseline over weeks 1 to 12 in monthly migraine days</b>			
Dose	100 mg (n = 222)	300 mg (n = 222)	Placebo (n = 222)
Days (SD or SE)	3.9 (NR)	4.3 (NR)	3.2 (NR)
Difference vs. PL, days (95% CI)	NR	NR	
P value	0.0179	0.0001	
Eptinezumab	Phase II Trial <sup>7</sup>		
<b>Change from baseline to weeks 5 to 8 in the frequency of migraine days</b>			
Dose		1,000 mg (n = 73)	Placebo (n = 78)
Days (SD)		5.6 (3.0)	4.6 (3.6)
Difference vs. PL, days (95% CI)		1.0 ( 2.1 to 0.1)	
P value		0.0306	
Erenumab	Phase III Trial (ARISE) <sup>16,32</sup>		
<b>Change from baseline to weeks 9 to 12 in monthly migraine days</b>			
Dose	70 mg (n = 289)		Placebo (n = 291)
Days (SD or SE)	2.9 (NR)		1.8 (NR)
Difference vs. PL, days (95% CI)	NR		

<i>P</i> value	< 0.001		
<b>Erenumab</b>	<b>Phase III Trial (STRIVE)<sup>17,18,32,33</sup></b>		
<b>Change from baseline in monthly migraine days between weeks 13 to 24</b>			
Dose	70 mg (n = 312)	140 mg (n = 318)	Placebo (n = 316)
Days (SE)	3.2 (0.2)	3.7 (0.2)	1.8 (0.2)
Difference vs. PL, days (95% CI)	1.4 ( 1.9 to 0.9)	1.9 ( 2.3 to 1.4)	
<i>P</i> value	< 0.001	< 0.001	
<b>Erenumab</b>	<b>Phase II Trial<sup>35</sup></b>		
<b>Change from baseline in monthly migraine days between weeks 9 to 12</b>			
Dose	70 mg (n = 107)		Placebo (n = 160)
Days (SE)	3.4 (0.4)		2.28 (0.3)
Difference vs. PL, days (95% CI)	1.1 ( 2.1 to 0.2)		
<i>P</i> value	0.021		
<b>Fremanezumab</b>	<b>Phase III Trial (HALO EM)<sup>21</sup></b>		
<b>Change from baseline in the monthly migraine days during the 12-week period</b>			
Dose	225 mg (n = NR)	675 mg (n = NR)	Placebo (n = NR)
Days (SD or SE)	3.7 (NR)	3.4 (NR)	2.2 (NR)
Difference vs. PL, days (95% CI)	NR	NR	
<i>P</i> value	< 0.0001	< 0.0001	
<b>Fremanezumab</b>	<b>Phase IIb Trial<sup>22</sup></b>		
<b>Change from baseline in the number of migraine days during weeks 9 to 12</b>			
Dose	225 mg (n = 95)	675 mg (n = 95)	Placebo (n = 104)
Days (SD)	6.27 (5.38)	6.09 (5.22)	3.46 (5.4)
Difference vs. PL, days (95% CI)	2.81 ( 4.07 to 1.55)	2.64 ( 3.90 to 1.38)	
<i>P</i> value	< 0.0001	< 0.0001	
<b>Galcanezumab</b>	<b>Phase III (EVOLVE-1)<sup>19</sup></b>		
<b>Change from baseline in monthly migraine days over 24 weeks</b>			
Dose	120 mg (n = NR)	240 mg (n = NR)	Placebo (n = NR)
Days (SD or SE)	4.7 (NR)	4.6 (NR)	2.8 (NR)
Difference vs. PL, days (95% CI)	NR	NR	
<i>P</i> value	< 0.001	< 0.001	
<b>Galcanezumab</b>	<b>Phase III (EVOLVE-2)<sup>19</sup></b>		
<b>Change from baseline in monthly migraine days over 24 weeks</b>			
Dose	120 mg (n = NR)	240 mg (n = NR)	Placebo (n = NR)
Days (SD or SE)	4.3 (NR)	4.2 (NR)	2.3
Difference vs. PL, days (95% CI)	NR	NR	
<i>P</i> value	< 0.001	< 0.001	
<b>Galcanezumab</b>	<b>Phase II<sup>38</sup></b>		
<b>Change from baseline to weeks 9 to 12 in monthly migraine days</b>			
Dose	150 mg (n = 108)		Placebo (n = 110)
Days (SD)	4.2 (3.1)		3.0 (3.0)
Difference vs. PL, days (90% CI)	1.2 ( 1.9 to 0.6)		
<i>P</i> value	0.003		

CI = confidence interval; NR = not reported; PL = placebo; SD = standard deviation; SE = standard error; vs. = versus.

<sup>a</sup> For detailed dosing information, please consult Table 2.

**Chronic Migraine:** Various primary outcomes were measured in the trials for chronic migraine.

*Eptinezumab:* With eptinezumab 100 mg and eptinezumab 300 mg, 31% and 33% of patients achieved a 75% reduction or more in migraine days over the course of the 12-week trial, respectively, compared with 21% of placebo patients ( $P < 0.05$  for both comparisons).

*Erenumab*: In a phase II trial, the mean difference in monthly migraine days was 2.5 days (95% CI, 3.5 to 1.4,  $P < 0.0001$ ) for both erenumab 70 mg and erenumab 140 mg compared with placebo.

*Fremanezumab* : In HALO-CM, the mean difference in monthly headache days was -2.1 days (SE 0.3) for the fremanezumab 675 mg initial dose followed by 225 mg group and -1.8 days (SE 0.3) for fremanezumab 675 mg single dose, compared with placebo ( $P < 0.001$  for both comparisons).

In a phase III trial, the mean difference in headache-hours during weeks 9 to 12 was -22.74 hours (95% CI, -44.28 to -1.21,  $P = 0.0386$ ) for fremanezumab 675 mg initial dose followed by 225 mg group and -30.41 hours (95% CI: -55.88 to -8.95,  $P = 0.0057$ ) for fremanezumab 900 mg, compared with placebo.

*Galcanezumab*: In REGAIN, the mean differences in monthly migraine days compared with placebo were 2.1 days ( $P = 0.001$ ) and 1.9 days ( $P = 0.001$ ) for galcanezumab 120 mg and galcanezumab 240 mg, respectively.

**Table 5: Change in Frequency of Migraines or Headaches<sup>a</sup> Chronic Migraine**

<b>Eptinezumab</b>		<b>Phase II Trial<sup>39</sup></b>		
<b>% patients achieving 75% reduction from baseline in migraine days over weeks 1 to 12</b>				
Dose	100 mg (n=NR)	300 mg (n = NR)	Placebo (n = NR)	
% patients (n/N)	31 (NR)	33 (NR)	21 (NR)	
Odds Ratio (95% CI)	NR	NR		
P value	< 0.05	< 0.05		
<b>Erenumab</b>		<b>Phase II Trial<sup>41</sup></b>		
<b>Change from baseline to weeks 9 to 12 in monthly migraine days</b>				
Dose	70 mg (n = 188)	140 mg (n = 187)	Placebo (n = 281)	
Days (SE)	6.6 (0.4)	6.6 (0.4)	4.2 (0.4)	
Difference vs. PL, days (95% CI)	2.5 ( 3.5 to 1.4)	2.5 ( 3.5 to 1.4)		
P value	< 0.0001	< 0.0001		
<b>Fremanezumab</b>		<b>Phase III Trial (HALO-CM)<sup>42,45</sup></b>		
<b>Change from baseline in monthly headache days over 3 months</b>				
Dose	675 mg/ 225 mg (n = 375)	675 mg/placebo (n = 375)	Placebo (n = 371)	
Days (SE)	4.6 (0.3)	4.3 (0.3)	2.5 (0.3)	
Difference vs. PL, days (95% CI)	2.1 (0.3)	1.8 (0.3)		
P value	< 0.001	< 0.001		
<b>Fremanezumab</b>		<b>Phase II Trial<sup>23</sup></b>		
<b>Change from baseline in headache-hours during weeks 9 to 12</b>				
Dose	675 mg/ 225 mg (n = 88)	900 mg (n = 87)	Placebo (n = 89)	
Hours (SD)	59.84 (80.38)	67.51 (79.37)	37.10 (79.44)	
Difference vs. PL, hours (95% CI)	22.74 ( 44.28 to 1.21)	30.41 ( 51.88 to 8.95)		
P value	0.0386	0.0057		
<b>Galcanezumab</b>		<b>Phase III (REGAIN)<sup>19,30</sup></b>		
<b>Change from baseline in monthly migraine days over 12 weeks</b>				
Dose	120 mg (n = 278)	240 mg (n = 277)	Placebo (n = 558)	
Days (SD or SE)	4.8 (NR)	4.6 (NR)	2.7 (NR)	
Difference vs. PL, days (95% CI)	NR	NR		
P value	0.001	0.001		

CI = confidence interval; NR = not reported; PL = placebo; SD = standard deviation; SE = standard error.

<sup>a</sup> For detailed dosing information, please consult Table 3.

## Safety

There was insufficient safety information in the phase III randomized controlled trials reported as conference abstracts or press releases and, therefore, only the safety results for published phase II and phase IIb trials are reported in Table 5.

No deaths were reported in the phase II and phase IIb trials.

The frequency of adverse events ranged from 44% to 72% with the anti-CGRP mAbs and from 39% to 67% with placebo. Adverse events most frequently reported included upper respiratory tract infection, nasopharyngitis, urinary tract infection, fatigue, back pain, muscle spasm, arthralgia, and abdominal pain, with similar frequencies across active and placebo arms. Injection site pain and injection site reactions were more frequent in the active groups compared with placebo.

A total of 1% to 2% of patients in the active treatment arms experienced a serious adverse event. These included:

- With eptinezumab 1,000 mg, one patient experienced chest pain, transient ischemic attack, conversion disorder, and dyspnea, and one patient had pyelonephritis requiring hospitalization, compared with one patient fracturing a fibula in the placebo group.
- With erenumab 70 mg, six patients experienced vertigo and migraine, intervertebral disc protrusion, appendicitis, costochondritis, fibroma, non-cardiac chest pain, and a fracture of the radius, whereas with erenumab 140 mg, two patients reported abdominal adhesions, abdominal pain, or cartilage injury. With placebo, seven patients experienced a serious adverse event, which included intervertebral disc protrusion, cholecystitis, migraine, pancreatitis, parotitis, urinary tract infection, and vomiting.
- With fremanezumab 225 mg, one patient fractured a fibula and one patient had a migraine associated with a hypertensive crisis, whereas with the 675 mg dose, one patient reported tremor and another reported antiphospholipid syndrome. In the group receiving an initial dose of 675 mg followed by fremanezumab 225 mg, one patient had pneumonia. At the higher dose of 900 mg, depression with suicide attempt, suicide attempt, and severe irritable bowel syndrome were reported in two patients. In the placebo group, one patient had nephrolithiasis.
- Details regarding the serious adverse events experienced with galcanezumab were not provided in the publications.

Withdrawals due to adverse events were infrequent. Reasons for stopping treatment because of adverse events included temperature intolerance, headache and migraine (erenumab 70 mg, all in one patient), fibula fracture and migraine associated with hypertensive crisis (fremanezumab 225 mg, one patient), and antiphospholipid syndrome and tremor (fremanezumab 675 mg, one patient). Two patients stopped erenumab 140 mg because of adverse events (constipation, fatigue, and metrorrhagia). Withdrawals because of an adverse event were reported in four patients and three patients administered with fremanezumab 675mg/225 mg and fremanezumab 900 mg, respectively; no reasons were provided in the publication.

Of note, CGRP neuropeptides, which are widely distributed in the body, are potent vasodilators. Thus, inhibiting CGRPs could potentially cause serious adverse events, including ischemic events.<sup>3,30</sup> So far, the trials showed no evidence of serious cardiovascular adverse events.

## Study Limitations

A common theme for all the trials is that the study population included a high proportion of women compared with men. This is expected considering that migraine headaches affect three times more women than men.<sup>2,4,5</sup> This raises concerns about the generalizability of the results to the male population. The majority of the study population was white, which also raises the concern about the generalizability of these study results in patients of other ethnic origins as medication efficacy and side effects can vary with ethnicity.<sup>46</sup>

The evidence on efficacy obtained so far for all four drugs is from placebo-controlled trials. There is a need to assess the efficacy of these drugs in comparison to standard of care in head-to-head trials in order to better determine their place in therapy.

Finally, some of the data were obtained from conference abstracts and press releases. There may be gaps in the results.

**Table 6: Harms from Published Phase II and Phase II/b Randomized Controlled Trials Number of Patients With Events**

EPTINEZUMAB					
	Episodic <sup>7</sup>				
	1,000 mg (n = 81)	Placebo (n = 82)			
<b>NoP (%): Adverse Events</b>	46 (57)	43 (52)			
<b>NoP (%): Serious Adverse Events</b>	2 (2)	1 (1)			
<b>NoP (%): Withdrawals Due to Adverse Events</b>	0	0			
ERENUMAB					
	Episodic <sup>35</sup>		Chronic <sup>41</sup>		
	70 mg (n = 106)	Placebo (n = 153)	140 mg (n = 188)	70 mg (n = 190)	Placebo (n = 282)
<b>NoP (%): Adverse Events</b>	54 (51)	80 (53)	88 (47)	83 (44)	110 (39)
<b>NoP (%): Serious Adverse Events</b>	1 (1)	0	2 (1)	6 (3)	7 (2)

<b>NoP (%): Withdrawals Due to Adverse Events</b>	1(1)	1 (< 1)	2(1)	0	2 (< 1)	
<b>FREMANEZUMAB</b>						
	<b>Episodic<sup>22</sup></b>			<b>Chronic<sup>23</sup></b>		
	<b>225 mg (n = 96)</b>	<b>675 mg (n = 96)</b>	<b>Placebo (n = 104)</b>	<b>675/225 mg (n = 88)</b>	<b>900 mg (n = 86)</b>	<b>Placebo (n = 89)</b>
<b>NoP (%): Adverse Events</b>	44 (46)	57 (59)	58 (56)	47 (53)	41 (48)	36 (40)
<b>NoP (%): Serious Adverse Events</b>	2 (2)	2 (2)	0	1 (1)	2 (2)	1 (1)
<b>NoP (%): Withdrawals Due to Adverse Events</b>	1 (1)	1 (1)	0	4 (5)	3 (3)	1 (1)
<b>GALCANEZUMAB</b>						
	<b>Episodic<sup>38</sup></b>					
	<b>150 mg (n = 107)</b>	<b>Placebo (n = 110)</b>				
<b>NoP (%): Adverse Events</b>	77 (72)	74 (67)				
<b>NoP (%): Serious Adverse Events</b>	2 (2)	4 (4)				
<b>NoP (%): Withdrawals Due to Adverse Events</b>	0	1 (1)				

NoP= number of patients.

## Concurrent Developments

PROMISE-2 is a phase III, randomized, double-blind, placebo- controlled trial of eptinezumab currently being conducted in the prevention of chronic migraines. The final data collection date for the primary outcomes measure (change in number of migraine days) is set for June 2018.<sup>47</sup>

Galcanezumab and fremanezumab are currently in phase III trials for the management of cluster headaches.<sup>48-53</sup>

Atogepant (AGN-241689, formerly MK-8031; Allergan) is currently being studied for episodic migraine prevention.<sup>54,55</sup>

Ubrogepant (MK-1602-006; Allergan) and rimegepant (BHV-3000; Biohaven Pharmaceuticals) are currently in phase III trials to evaluate their efficacy in the treatment of acute migraines.<sup>54</sup>

Lasmiditan (COL-144 or LY573144; Eli Lilly.), a 5-HT<sub>1F</sub> receptor agonist (commonly known as a “ditan”) is currently in phase III clinical trials for the treatment of acute migraine.<sup>54,56,57 58</sup>

## Implementation Issues

Access to anti-CGRP mAbs is currently limited, as they have not been approved in any country. Therefore none of the drugs are currently marketed in Canada. Although the cost of these drugs has not been determined, they will potentially have a significant budget impact because of the high cost of biologics and the prevalence of episodic and chronic migraines.

Eptinezumab is the only drug administered intravenously. Intravenous administration could pose a challenge depending on the accessibility and availability of treatment centres. Eptinezumab and fremanezumab are administered every three months; this administration schedule could be desirable to patients who prefer less-frequent dosing regimens.

The appropriate duration of treatment and whether these drugs can eventually be tapered down or discontinued still needs to be clarified. Information on how these drugs will be used with other preventive migraine therapies is lacking.<sup>30</sup>

## References

1. Negro A, Curto M, Lionetto L, Giamberardino MA, Martelletti P. Chronic migraine treatment: from OnabotulinumtoxinA onwards. *Expert Rev Neurother*. 2016 Oct;16(10):1217-27.
2. Mitsikostas DD, Rapoport AM. New players in the preventive treatment of migraine. *BMC Med* [Internet]. 2015 Nov 10 [cited 2017 Dec 18];13:279. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4641418>
3. Pellesi L, Guerzoni S, Pini LA. Spotlight on anti-CGRP monoclonal antibodies in migraine: the clinical evidence to date. *Clin Pharmacol Drug Dev*. 2017 Apr 14.
4. Robinson G. Identifying and treating chronic migraine. *Can J Diabetes*. 2015 Oct;43-6.
5. Mitsikostas DD, Reuter U. Calcitonin gene-related peptide monoclonal antibodies for migraine prevention: comparisons

- across randomized controlled studies. *Curr Opin Neurol*. 2017 Jun;30(3):272-80.
6. Bigal ME, Walter S, Rapoport AM. Therapeutic antibodies against CGRP or its receptor. *Br J Clin Pharmacol* [Internet]. 2015 Jun [cited 2017 Dec 18];79(6):886-95. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4456121>
  7. Dodick DW, Goadsby PJ, Silberstein SD, Lipton RB, Olesen J, Ashina M, et al. Safety and efficacy of ALD403, an antibody to calcitonin gene-related peptide, for the prevention of frequent episodic migraine: a randomised, double-blind, placebo-controlled, exploratory phase 2 trial. *Lancet Neurol*. 2014 Nov;13(11):1100-7.
  8. Tso AR, Goadsby PJ. Anti-CGRP monoclonal antibodies: the next era of migraine prevention? *Curr Treat Options Neurol* [Internet]. 2017 Aug [cited 2017 Aug 25];19(8):27. Available from: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5486583/pdf/11940\\_2017\\_Article\\_463.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5486583/pdf/11940_2017_Article_463.pdf)
  9. Edvinsson L, Ho TW. CGRP receptor antagonism and migraine. *Neurotherapeutics* [Internet]. 2010 Apr [cited 2017 Oct 16];7(2):164-75. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5084097>
  10. Alder BioPharmaceuticals. Bothell (WA): Alder BioPharmaceuticals. Press release, Alder BioPharmaceuticals announces positive eptinezumab phase 3 results for prevention of frequent episodic migraine; 2017 Jun 27
  11. Drug and health product submissions under review (SUR). 2017 Nov 7 [cited 2017 Dec 1]. In: Drug and health product review and approval [Internet]. Ottawa (ON): Government of Canada. Available from: [https://www.canada.ca/en/health-canada/services/drug-health-product-review-approval/submissions-under-review.html?\\_ga=2.114239301.378295653.1512062912-591177382.1389892045](https://www.canada.ca/en/health-canada/services/drug-health-product-review-approval/submissions-under-review.html?_ga=2.114239301.378295653.1512062912-591177382.1389892045).
  12. Novartis [Internet]. Basel (CH): Novartis. Press release, Novartis achieves important regulatory milestone for AMG 334 (erenumab) in migraine prevention with EMA filing acceptance; 2017 Jun 21 [cited 2017 Sep 1]. Available from: <https://www.novartis.com/news/media-releases/novartis-achieves-important-regulatory-milestone-amg-334-erenumab-migraine>
  13. Cision PR Newswire [Internet]. New York: Cision PR Newswire. FDA accepts biologics license application for Aimovig™ (erenumab); 2017 Jul 20 [cited 2017 Sep 13]. Available from: <http://www.prnewswire.com/news-releases/fda-accepts-biologics-license-application-for-aimovig-erenumab-300491861.html>
  14. Hee DH. BLA submitted for migraine prevention drug fremanezumab. In: The Right Dose of Information [Internet]. London: Haymarket Media, Inc.; 2017 Oct 17 [cited 2017 Oct 23]. Available from: <http://www.empr.com/drugs-in-the-pipeline/fremanezumab-migraine-prevention-biologics-license-application/article/700664/>
  15. PharmaTimes [Internet]. London: PharmaTimes. Lilly files anti-CGRP drug for migraine; 2017 Dec 12 [cited 2017 Dec 18]. Available from: [http://www.pharmatimes.com/news/lilly\\_files\\_anti\\_cgrp\\_drug\\_for\\_migraine\\_1214101?utm\\_source=PT%20Weekly%20Newsletter&utm\\_medium=email&utm\\_campaign=pt%20weekly%20news%20alert&utm\\_source=PT%20Weekly%20Newsletter&utm\\_medium=email&utm\\_campaign=pt%20weekly%20news%20alert&](http://www.pharmatimes.com/news/lilly_files_anti_cgrp_drug_for_migraine_1214101?utm_source=PT%20Weekly%20Newsletter&utm_medium=email&utm_campaign=pt%20weekly%20news%20alert&utm_source=PT%20Weekly%20Newsletter&utm_medium=email&utm_campaign=pt%20weekly%20news%20alert&)
  16. Dodick DW, Ashina M, Kudrow D, Lanteri-Minet M, Osipova V, Palmer K, et al. A phase 3, randomized, double-blind, placebo-controlled study to evaluate the efficacy and safety of erenumab in migraine prevention: primary results of the ARISE trial. *Headache*. 2017 Jun;191-2.
  17. Goadsby PJ, Reuter U, Bonner J, Broessner G, Hallstrom Y, Zhang F, et al. Phase 3, randomized, double-blind, placebo-controlled study to evaluate the efficacy and safety of erenumab (AMG 334) in migraine prevention: primary results of the STRIVE Trial. *Eur J Neurol*. 2017;117.
  18. Goadsby PJ, Reuter U, Bonner J, Broessner G, Hallstrom Y, Zhang F, et al. Phase 3, randomized, double-blind, placebo-controlled study to evaluate the efficacy and safety of erenumab (AMG 334) in migraine prevention: primary results of the STRIVE trial. *Headache*. 2017;128-9.
  19. Eli Lilly [Internet]. Indianapolis (IN): Eli Lilly and Company. Press release, Lilly announces positive results for three phase 3 studies of galcanezumab for the prevention of episodic and chronic migraine; 2017 May 12 [cited 2017 Sep 19]. Available from: <https://investor.lilly.com/releasedetail.cfm?releaseid=1026201>
  20. Pharmiweb.com [Internet]. Bracknell (GB): Pharmiweb.com. Press release, Teva to present fremanezumab data on migraine prevention at the 18th Congress of the International Headache Society; 2017 Sep 5 [cited 2017 Sep 19]. Available from: [http://www.pharmiweb.com/pressreleases/pressrel.asp?ROW\\_ID=241311#.WcFeUGfbJeX](http://www.pharmiweb.com/pressreleases/pressrel.asp?ROW_ID=241311#.WcFeUGfbJeX)
  21. Teva Pharmaceutical Industries Ltd. [Internet]. Petach Tikva (IL): Teva Pharmaceutical Industries Ltd. Press release, Teva's fremanezumab meets all primary & secondary endpoints across both monthly and quarterly dosing regimens in phase III study in episodic migraine prevention; 2017 [cited 2017 Sep 19]. Available from: [http://www.tevapharm.com/news/teva\\_s\\_fremanezumab\\_meets\\_all\\_primary\\_secondary\\_endpoints\\_across\\_both\\_monthly\\_and\\_quarterly\\_dosing\\_regimens\\_in\\_phase\\_iii\\_study\\_in\\_episodic\\_migraine\\_prevention\\_06\\_17.aspx](http://www.tevapharm.com/news/teva_s_fremanezumab_meets_all_primary_secondary_endpoints_across_both_monthly_and_quarterly_dosing_regimens_in_phase_iii_study_in_episodic_migraine_prevention_06_17.aspx)
  22. Bigal ME, Dodick DW, Rapoport AM, Silberstein SD, Ma Y, Yang R, et al. Safety, tolerability, and efficacy of TEV-48125 for preventive treatment of high-frequency episodic migraine: a multicentre, randomised, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol*. 2015 Nov;14(11):1081-90.
  23. Bigal ME, Edvinsson L, Rapoport AM, Lipton RB, Spierings EL, Diener HC, et al. Safety, tolerability, and efficacy of TEV-48125 for preventive treatment of chronic migraine: a multicentre, randomised, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol*. 2015 Nov;14(11):1091-100.
  24. Sun-Edelstein C, Rapoport AM. Update on the pharmacological treatment of chronic migraine. *Curr Pain Headache Rep*. 2016 Jan;20(1):6.
  25. Novartis [Internet]. Basel (CH): Novartis International AG. Press release, Novartis presents new analysis demonstrating AMG 334 (erenumab) significantly reduced monthly migraine days in patients who failed previous preventive therapies; 2017 Sep 7 [cited 2017 Sep 19]. Available from: <https://www.novartis.com/news/media-releases/novartis-presents-new-analysis-demonstrating-amg-334-erenumab-significantly>
  26. Roceanu A, Antochi F, Bajenaru O. New molecules in migraine treatment. *Farmacia*. 2015;63(4):475-81.
  27. Topamax (topiramate): topiramate tablets, House Std.25, 100 and 200 mg; topiramate sprinkle capsules, House Std.15 and 25 mg [product monograph] [Internet]. Toronto (ON): Janssen Inc.; 2017 Apr 28. [cited 2017 Dec 12]. Available from: [http://www.janssen.com/canada/sites/www\\_janssen\\_com\\_canada/files/prod\\_files/live/topamax\\_cpm.pdf](http://www.janssen.com/canada/sites/www_janssen_com_canada/files/prod_files/live/topamax_cpm.pdf)
  28. Tarighat Esfanjani A, Mahdavi R, Ebrahimi Mameghani M, Talebi M, Nikniaz Z, Safaiyan A. The effects of magnesium, L-carnitine, and concurrent magnesium-L-carnitine supplementation in migraine prophylaxis. *Biol Trace Elem Res*. 2012 Dec;150(1-3):42-8.
  29. Alder Biopharmaceuticals, Inc. A multicenter assessment of ALD403 in frequent episodic migraine (PROMISE 1). 2017

- Apr 11 [cited 2017 Dec 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02559895?term=ALD+403&rank=6> NLM Identifier: NCT02559895.
30. Keller DM. CGRP monoclonals: safe, effective for migraine in phase 3 trials. In: Medscape [Internet]. New York (NY): Medscape LLC; 2017 Sep 22 [cited 2017 Oct 4]. Available from: <https://www.medscape.com/viewarticle/886068>
  31. Amgen. Study to evaluate the efficacy and safety of AMG 334 compared to placebo in migraine prevention (ARISE). 2017 May 30 [cited 2017 Sep 13]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02483585> NLM Identifier: NCT02483585.
  32. Brauser D. Phase 3 STRIVE and ARISE trials show efficacy, safety for erenumab in migraine prevention [Internet]. New York (NY): Medscape LLC; 2017 Apr 26. [cited 2017 Dec 1]. Available from: <https://www.medscape.com/viewarticle/879161>
  33. Goadsby PJ, Reuter U, Hallstrom Y, Broessner G, Bonner JH, Zhang F, et al. A controlled trial of erenumab for episodic migraine. *N Engl J Med*. 2017 Nov 30;377(22):2123-32.
  34. Amgen. Study to evaluate the efficacy and safety of AMG 334 in migraine prevention (STRIVE). 2017 Jun 20 [cited 2017 Sep 13]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02456740> NLM Identifier: NCT02456740.
  35. Sun H, Dodick DW, Silberstein S, Goadsby PJ, Reuter U, Ashina M, et al. Safety and efficacy of AMG 334 for prevention of episodic migraine: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Neurol*. 2016 Apr;15(4):382-90.
  36. Eli Lilly and Company. Evaluation of LY2951742 in the prevention of episodic migraine- the EVOLVE-1 study (EVOLVE-1). 2017 Apr 19 [cited 2017 Sep 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02614183?term=LY+2951742&draw=1&rank=7> NLM Identifier: NCT02614183.
  37. Eli Lilly and Company. Evaluation of LY2951742 in the prevention of episodic migraine- the EVOLVE-2 study (EVOLVE-2). 2017 Apr 19 [cited 2017 Dec 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02614196?term=LY+2951742&draw=1&rank=6> NLM Identifier: NCT02614196.
  38. Dodick DW, Goadsby PJ, Spierings EL, Scherer JC, Sweeney SP, Grayzel DS. Safety and efficacy of LY2951742, a monoclonal antibody to calcitonin gene-related peptide, for the prevention of migraine: a phase 2, randomised, double-blind, placebo-controlled study. *Lancet Neurol*. 2014 Sep;13(9):885-92.
  39. Smith J, Dodick DW, Goadsby PJ, Silberstein SD, Lipton RB, Hirman J. Randomized, double-blind, placebo-controlled trial of ALD403 (eptinezumab), an anti-CGRP monoclonal antibody for the prevention of chronic migraine. *Headache*. 2017;57 Suppl 3:130.
  40. Alder Biopharmaceuticals, Inc. A multicenter assessment of ALD403 in chronic migraine. 2017 Feb 17 [cited 2017 Sep 13]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02275117> NLM Identifier: NCT02275117.
  41. Tepper S, Ashina M, Reuter U, Brandes JL, Dolezil D, Silberstein S, et al. Safety and efficacy of erenumab for preventive treatment of chronic migraine: a randomised, double-blind, placebo-controlled phase 2 trial. *Lancet Neurol*. 2017 Jun;16(6):425-34.
  42. Silberstein SD, Dodick DW, Bigal ME, Yeung PP, Goadsby PJ, Blankenbiller T, et al. Fremanezumab for the preventive treatment of chronic migraine. *N Engl J Med*. 2017 Nov 30;377(22):2113-22.
  43. Teva Branded Pharmaceutical Products, R&D Inc. Comparing efficacy and safety of 2 dose regimens of subcutaneous administration of TEV-48125 versus placebo for the preventive treatment of chronic migraine. 2017 Aug 11 [cited 2017 Sep 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02621931?term=TEV-48125&rank=3> NLM Identifier: NCT02621931.
  44. Eli Lilly and Company. Evaluation of LY2951742 in the prevention of chronic migraine (REGAIN). 2015 Nov 23 [cited 2017 Sep 29; Last update posted: 2017 Apr 20]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://www.clinicaltrials.gov/ct2/show/NCT02614261?cond=LY2951742&draw=1&rank=3> Identifier: NCT02614261.
  45. Robinson R. Erenumab and fremanezumab found to reduce chronic migraine days. In: Neurology today conference reporter: American Headache Society Annual Meeting [blog on the Internet]. Minneapolis (MN): American Academy of Neurology; 2017 Jun 15 [cited 2017 Sep 13]. Available from: <http://journals.lww.com/neurotodayonline/blog/NeurologyTodayConferenceReporterAmericanHeadacheSocietyAnnualMeeting/pages/post.aspx?PostID=12>
  46. Yasuda SU, Zhang L, Huang SM. The role of ethnicity in variability in response to drugs: focus on clinical pharmacology studies. *Clin Pharmacol Ther*. 2008 Sep;84(3):417-23.
  47. Alder Biopharmaceuticals, Inc. Evaluation of ALD403 (eptinezumab) in the prevention of chronic migraine (PROMISE 2). 2017 Aug 11 [cited 2017 Dec 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02974153?term=ALD+403&rank=5> NLM Identifier: NCT02974153.
  48. Eli Lilly and Company. A study of LY2951742 in participants with episodic cluster headache. 2015 Mar 19 [cited 2017 Sep 29; Last update posted: 2017 Sep 26]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://www.clinicaltrials.gov/ct2/show/NCT02397473?cond=LY2951742&draw=1&rank=6> Identifier: NCT02397473.
  49. Eli Lilly and Company. A study of LY2951742 in participants with chronic cluster headache. 2015 May 6 [cited 2017 Sep 29; Last update posted: 2017 Sep 26]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://www.clinicaltrials.gov/ct2/show/NCT02438826?cond=LY2951742&draw=2&rank=11> Identifier: NCT02438826.
  50. Eli Lilly and Company. A study of LY2951742 (galcanezumab) in participants with cluster headache. 2016 Jun 9 [cited 2017 Sep 29; Last update posted: 2017 Jun 27]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://www.clinicaltrials.gov/ct2/show/NCT02797951?cond=LY2951742&draw=1&rank=12> Identifier: NCT02797951.

51. Teva Branded Pharmaceutical Products, R&D Inc. A study to evaluate the efficacy and safety of TEV-48125 (fremanezumab) for the prevention of episodic cluster headache (ECH). 2016 Oct 25 [cited 2017 Sep 29; Last update posted: 2017 Aug 24]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02945046?cond=TEV48125&rank=7> Identifier: NCT02945046.
52. Teva Branded Pharmaceutical Products, R&D Inc. A study comparing the efficacy and safety of TEV-48125 (fremanezumab) for the prevention of chronic cluster headache (CCH). 2016 Nov 11 [cited 2017 Sep 29; Last update posted: 2017 Jul 19]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT02964338?cond=TEV48125&rank=6> Identifier: NCT02964338.
53. Teva Branded Pharmaceutical Products, R&D Inc. A study to explore the long-term safety of TEV-48125 (fremanezumab) for the prevention of cluster headache (ENFORCE). 2017 Mar 23 [cited 2017 Sep 29; Last update posted: 2017 Sep 20]. In: ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2000 - . Available from: <https://clinicaltrials.gov/ct2/show/NCT03107052?cond=TEV48125&rank=1> Identifier: NCT03107052.
54. Schuster NM, Rapoport AM. Calcitonin gene-related peptide-targeted therapies for migraine and cluster headache: a review. *Clin Neuropharmacol*. 2017 Jul;40(4):169-74.
55. Biohaven Pharmaceuticals [Internet]. New Haven (CT): Biohaven Pharmaceuticals. News release, Biohaven enrolls first patient in pivotal phase 3 clinical trial of oral rimegepant for the acute treatment of migraine; 2017 Jul 27 [cited 2017 Sep 13]. Available from: <http://biohavenpharma.com/biohaven-enrolls-first-patient-in-pivotal-phase-3-clinical-trial-of-oral-rimegepant-for-the-acute-treatment-of-migraine/>
56. Schuster NM, Vollbracht S, Rapoport AM. Emerging treatments for the primary headache disorders. *Neurol Sci*. 2015 May;36 Suppl 1:109-13.
57. Lilly [Internet]. Indianapolis (IN): Lilly. News release, Lilly announces positive results for second phase 3 study of lasmiditan for the acute treatment of migraine; 2017 [cited 2017 Sep 13]. Available from: <https://investor.lilly.com/releasedetail.cfm?ReleaseID=1036101>
58. Lilly [Internet]. Indianapolis (IN): Lilly. News release, Lilly and CoLucid Pharmaceuticals announce agreement for Lilly to acquire CoLucid; 2017 [cited 2017 Dec 12]. Available from: <https://investor.lilly.com/releaseDetail.cfm?ReleaseID=1008267>

## Appendix 1: Small-Molecule CGRP Receptor Antagonists to Treat Acute Migraine No Longer in Development

Drug	Development Name	Development Status
Olcegepant	BIBN4096BS	Did not proceed beyond phase II because of modest efficacy
Telcagepant	MK-0974	Six completed phase III trials; sponsor has stopped clinical development because of liver toxicity
NA	BI 44370 TA	Development phase discontinued; reasons unknown
NA	MK-3207	Development discontinued because of liver toxicity

NA = not applicable.

## About This Document

Authors: Abas Ibekwe, BSc, Pharm D Candidate; Christine Perras, MPH; Monika Mierzwinski-Urban, MLIS

Cite As: Monoclonal antibodies to prevent migraine headaches. Ottawa: CADTH; 2018 Feb. (CADTH issues in emerging health technologies, issue 167).

Acknowledgments: CADTH thanks the external reviewers who kindly provided comments on an earlier draft of this bulletin.

Disclaimer: The information in this document is intended to help Canadian health care decision-makers, health care professionals, health systems leaders, and policymakers make well-informed decisions and thereby improve the quality of health care services. While patients and others may access this document, the document is made available for informational purposes only and no representations or warranties are made with respect to its fitness for any particular purpose. The information in this document should not be used as a substitute for professional medical advice or as a substitute for the application of clinical judgment in respect of the care of a particular patient or other professional judgment in any decision-making process. The Canadian Agency for Drugs and Technologies in Health (CADTH) does not endorse any information, drugs, therapies, treatments, products, processes, or services.

While care has been taken to ensure that the information prepared by CADTH in this document is accurate, complete, and up-to-date as at the applicable date the material was first published by CADTH, CADTH does not make any guarantees to that effect. CADTH does not guarantee and is not responsible for the quality, currency, propriety, accuracy, or reasonableness of any statements, information, or conclusions contained in any third-party materials used in preparing this document. The views and opinions of third parties published in this document do not necessarily state or reflect those of CADTH.

CADTH is not responsible for any errors, omissions, injury, loss, or damage arising from or relating to the use (or misuse) of any

information, statements, or conclusions contained in or implied by the contents of this document or any of the source materials.

This document may contain links to third-party websites. CADTH does not have control over the content of such sites. Use of third-party sites is governed by the third-party website owners' own terms and conditions set out for such sites. CADTH does not make any guarantee with respect to any information contained on such third-party sites and CADTH is not responsible for any injury, loss, or damage suffered as a result of using such third-party sites. CADTH has no responsibility for the collection, use, and disclosure of personal information by third-party sites.

Subject to the aforementioned limitations, the views expressed herein are those of CADTH and do not necessarily represent the views of Canada's federal, provincial, or territorial governments or any third party supplier of information.

This document is prepared and intended for use in the context of the Canadian health care system. The use of this document outside of Canada is done so at the user's own risk.

This disclaimer and any questions or matters of any nature arising from or relating to the content or use (or misuse) of this document will be governed by and interpreted in accordance with the laws of the Province of Ontario and the laws of Canada applicable therein, and all proceedings shall be subject to the exclusive jurisdiction of the courts of the Province of Ontario, Canada.

The copyright and other intellectual property rights in this document are owned by CADTH and its licensors. These rights are protected by the Canadian Copyright Act and other national and international laws and agreements. Users are permitted to make copies of this document for non-commercial purposes only, provided it is not modified when reproduced and appropriate credit is given to CADTH and its licensors.

About CADTH: CADTH is an independent, not-for-profit organization responsible for providing Canada's health care decision-makers with objective evidence to help make informed decisions about the optimal use of drugs, medical devices, diagnostics, and procedures in our health care system.

Funding: CADTH receives funding from Canada's federal, provincial, and territorial governments, with the exception of Quebec.

Contact [requests@cadth.ca](mailto:requests@cadth.ca) with inquiries about this notice or legal matters relating to CADTH services.