

## CADTH OPTIMAL USE REPORT

# Interventions for the Treatment of Obstructive Sleep Apnea in Adults: Recommendations

Service Line:	Optimal Use
Issue Number:	volume 6, no. 1c
Publication Date:	March 2017
Report Length:	22 Pages

**Cite As:** Interventions for the treatment of obstructive sleep apnea in adults: recommendations. Ottawa: CADTH; Mar 2017. (CADTH Optimal Use Report; vol.6, no.1c).

**ISSN:** 1927-0127 (online)

**Disclaimer:** The information in this document is intended to help Canadian health care decision-makers, health care professionals, health systems leaders, and policy-makers make well-informed decisions and thereby improve the quality of health care services. While patients and others may access this document, the document is made available for informational purposes only and no representations or warranties are made with respect to its fitness for any particular purpose. The information in this document should not be used as a substitute for professional medical advice or as a substitute for the application of clinical judgment in respect of the care of a particular patient or other professional judgment in any decision-making process. The Canadian Agency for Drugs and Technologies in Health (CADTH) does not endorse any information, drugs, therapies, treatments, products, processes, or services.

While care has been taken to ensure that the information prepared by CADTH in this document is accurate, complete, and up-to-date as at the applicable date the material was first published by CADTH, CADTH does not make any guarantees to that effect. CADTH does not guarantee and is not responsible for the quality, currency, propriety, accuracy, or reasonableness of any statements, information, or conclusions contained in any third-party materials used in preparing this document. The views and opinions of third parties published in this document do not necessarily state or reflect those of CADTH.

CADTH is not responsible for any errors, omissions, injury, loss, or damage arising from or relating to the use (or misuse) of any information, statements, or conclusions contained in or implied by the contents of this document or any of the source materials.

This document may contain links to third-party websites. CADTH does not have control over the content of such sites. Use of third-party sites is governed by the third-party website owners' own terms and conditions set out for such sites. CADTH does not make any guarantee with respect to any information contained on such third-party sites and CADTH is not responsible for any injury, loss, or damage suffered as a result of using such third-party sites. CADTH has no responsibility for the collection, use, and disclosure of personal information by third-party sites.

Subject to the aforementioned limitations, the views expressed herein are those of CADTH and do not necessarily represent the views of Canada's federal, provincial, or territorial governments or any third party supplier of information.

This document is prepared and intended for use in the context of the Canadian health care system. The use of this document outside of Canada is done so at the user's own risk.

This disclaimer and any questions or matters of any nature arising from or relating to the content or use (or misuse) of this document will be governed by and interpreted in accordance with the laws of the Province of Ontario and the laws of Canada applicable therein, and all proceedings shall be subject to the exclusive jurisdiction of the courts of the Province of Ontario, Canada.

The copyright and other intellectual property rights in this document are owned by CADTH and its licensors. These rights are protected by the Canadian *Copyright Act* and other national and international laws and agreements. Users are permitted to make copies of this document for non-commercial purposes only, provided it is not modified when reproduced and appropriate credit is given to CADTH and its licensors.

**About CADTH:** CADTH is an independent, not-for-profit organization responsible for providing Canada's health care decision-makers with objective evidence to help make informed decisions about the optimal use of drugs, medical devices, diagnostics, and procedures in our health care system.

**Funding:** CADTH receives funding from Canada's federal, provincial, and territorial governments, with the exception of Quebec.

## Table of Contents

ABBREVIATIONS .....	4
SUMMARY OF RECOMMENDATIONS.....	5
Technology .....	6
Policy Question.....	6
METHODS .....	7
DETAILED RECOMMENDATIONS .....	7
Rationale .....	7
Considerations.....	8
BACKGROUND .....	10
Research questions .....	10
SUMMARY OF THE EVIDENCE .....	11
Clinical Evidence .....	11
Economic Evidence .....	12
Patient Perspectives and Experiences Evidence.....	12
Ethical Issues.....	13
Implementation Issues.....	14
Environmental Impact.....	14
RESEARCH GAPS .....	14
APPENDIX 1: HTERP .....	18
HTERP Core Members.....	18
Expert Members .....	18
Conflict of Interest.....	18
APPENDIX 2: HTERP DELIBERATIVE FRAMEWORK.....	19

## Abbreviations

AHI	apnea-hypopnea index
CPAP	continuous positive airway pressure
EDS	excessive daytime sleepiness
GTA	genial tubercle advancement
HTA	health technology assessment
HTERP	Health Technology Expert Review Panel
MAD	mandibular advancement device
MMA	maxillomandibular advancement
OSA	obstructive sleep apnea
PAP	positive airway pressure
QALY	quality-adjusted life year
QoL	quality of life

## Summary of Recommendations

Obstructive sleep apnea (OSA) is characterized by a narrowing and collapse of the upper airway during sleep.<sup>1,2</sup> The prevalence of OSA is reported to be 15% in males and 5% in females.<sup>3,4</sup> The major symptoms include snoring, unrefreshing sleep, excessive daytime sleepiness (EDS), lack of concentration, impaired memory, and lower quality of life.<sup>5,6</sup> Aging, the male sex, and obesity are the main risk factors for OSA.<sup>7,8</sup> Untreated OSA is associated with motor vehicle accidents, cardiovascular disease, stroke, hypertension, diabetes, cognitive dysfunction, and all-cause mortality.<sup>1,6,9-11</sup>

The goal of treatment of OSA is to reduce the apnea-hypopnea index (AHI) — an index used to indicate the severity of sleep apnea — increase blood oxygen levels, and improve cardiorespiratory indicators. The AHI measures the number of apnea or hypopnea events per hour.<sup>12</sup> Although continuous positive airway pressure (CPAP) is the standard for treating OSA, between 29% and 83% of patients do not comply with regular device use.<sup>13-15</sup> Adherence with oral appliances, including mandibular advancement devices (MADs), is not as well documented, but is regarded as superior to CPAP adherence.<sup>16,17</sup> Surgical interventions for OSA treatment are invasive procedures for which evidence of effectiveness and safety is unclear.<sup>12</sup> For patients with mild or asymptomatic OSA, lifestyle interventions such as exercise programs, diet changes, and positional therapies may be an option for treatment before proceeding to other interventions.<sup>18</sup>

Across jurisdictions, OSA is associated with a substantial economic and societal burden.<sup>7,8,19</sup> Currently, public coverage for treatment of OSA varies widely across Canada, with some provinces supporting CPAP therapy for OSA patients, but the criteria and type of reimbursement varies.<sup>2</sup> No provincial programs reimburse oral appliance costs while some federal programs do reimburse eligible patients.

Given the range of clinical presentation, symptoms, and severity, recommending the most appropriate treatment for OSA patients can be challenging. To facilitate decision-making, CADTH conducted a health technology assessment (HTA) on the clinical effectiveness and cost-effectiveness of interventions for the treatment of OSA in adults. Patient perspectives and experiences, ethical and implementation issues, and environmental factors related to therapy selection for OSA in adults were also considered in an evaluation of the appropriate use of OSA interventions.<sup>20</sup>

1. For patients with mild OSA who are overweight or obese, the Health Technology Expert Review Panel (HTERP) recommends lifestyle interventions. For patients with mild OSA who are not overweight or obese, HTERP does not recommend active treatment.
2. For patients with moderate or severe OSA, HTERP recommends continuous positive airway pressure (CPAP). For patients with moderate or severe OSA for whom CPAP is unacceptable, oral appliances are recommended.
3. HTERP does not recommend surgical maxillomandibular advancement in patients with OSA, unless other interventions have failed or are unacceptable to the patient.

## Technology

Treatment of obstructive sleep apnea (OSA) includes a wide range of options.<sup>21</sup>

Continuous positive airway pressure (CPAP) forces air into the upper airways to prevent soft tissues from collapsing and is considered the gold standard for the treatment of OSA.<sup>12,21,22</sup> Other positive airway pressure (PAP) technologies, such as autotitrating PAP and bilevel PAP, may be offered to patients with specific needs.<sup>12,21</sup> Another treatment option is nasal expiratory PAP valves, which are disposable devices that use a patient's own breathing to create positive end-expiratory pressure that prevents airway collapse.<sup>23</sup>

Oral appliances, including the mandibular advancement devices (MADs) and tongue-retaining devices (TRDs), can be offered as an alternative to CPAP.<sup>12,24,25</sup> For patients with mild or asymptomatic OSA, lifestyle interventions such as exercise programs, diet changes, and positional therapies may be proposed.<sup>18</sup>

Surgical maxillomandibular advancement (MMA) permanently pulls the lower jaw forward to create more space and prevent airway collapse.<sup>26,27</sup> Genial tubercle advancement (GTA) is a surgical intervention that removes bone tissue from the chin and pulls the base of the tongue forward to create more airway space, and can be performed in conjunction with MMA or other surgeries to potentially improve therapeutic success.<sup>28,29</sup>

## Policy Question

What is the optimal use of PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle interventions for treatment of adults with OSA?

## Methods

CADTH conducted a HTA to assess the clinical effectiveness, cost-effectiveness, patient perspectives and experiences, ethical issues, implementation issues, and environmental impact of PAP devices, oral appliances, surgical interventions, and lifestyle interventions for the treatment of OSA in adults.<sup>20</sup> The Health Technology Expert Review Panel (HTERP) (Appendix 1) developed recommendations about interventions for the treatment of OSA based on the evidence presented in the HTA report. HTERP members reviewed the evidence, discussed all elements of the HTERP deliberative framework,<sup>30</sup> and developed a consensus-based recommendation through discussion and deliberation. See Appendix 2 for details.

Additional information on the HTERP process is found on the HTERP page of the CADTH website: <https://www.cadth.ca/collaboration-and-outreach/advisory-bodies/health-technology-expert-review-panel>.

## Detailed Recommendations

The objective of these recommendations is to provide advice for Canadian health care decision-makers about the optimal use of interventions for the treatment of OSA of varying severity in adults. These recommendations are relevant for patients who were diagnosed with any severity of OSA and were either treatment-naïve or previously treated, as measured objectively by polysomnography or portable monitoring.

1. For patients with mild OSA who are overweight or obese, HTERP recommends lifestyle interventions. For patients with mild OSA who are not overweight or obese, HTERP does not recommend active treatment.
2. For patients with moderate or severe OSA, HTERP recommends CPAP. For patients with moderate or severe OSA for whom CPAP is unacceptable, oral appliances are recommended.
3. HTERP does not recommend surgical MMA in patients with OSA, unless other interventions have failed or are unacceptable to the patient.

## Rationale

The results of the clinical review indicate that, while various interventions may have similar and only marginal effects on improving sleepiness across mild-to-severe cases of OSA, CPAP may have the largest effect on improving OSA severity, if patients comply with the therapy. Further, the clinical and economic data indicate that patients with moderate OSA may benefit most from MADs, and those with severe OSA may benefit most from CPAP. More specifically, the results suggest that CPAP is more effective than MADs for EDS in adults with severe OSA. Although treatment is not

recommended in patients with mild OSA, the committee suggested that treatment may be considered in patients with mild OSA who are symptomatic.

Surgical MMA, with or without GTA, was the most clinically effective and cost-effective intervention at a willingness-to-pay threshold of \$17,125 per quality-adjusted life-year (QALY) in patients with very severe OSA. Unlike other interventions for OSA, this is an invasive procedure. In addition, the findings on MMA, with or without GTA, were obtained from small, uncontrolled pre- and post-treatment studies of highly selected patients, and therefore MMA is not recommended for most patients with OSA.

There were no major adverse events reported for most OSA interventions.

Some of the evidence was deemed to be of high quality, but there are quality concerns for some studies due to their eligibility criteria, sample sizes, and uncontrolled study design.

Patient factors that influence whether people seek and commence OSA treatment are individualized and contextual. People with OSA usually perform a trade-off between the benefits of OSA treatment and their discomfort with the intervention. All treatments had some degree of discomfort, and this discomfort may change over time as patients become accustomed to the device. There is also a recovery time for surgery. For some patients, especially those with mild OSA, these feelings of discomfort were enough of a deterrent that therapy was discontinued. For others, the physical, mental, and social benefits experienced from using an intervention for OSA were motivation to continue treatments. For those using CPAP, the sense of embarrassment and perceived unattractive appearance while using the device might be a reason for nonadherence. Those with supportive partners may be able to persevere and continue with treatment, though not all spouses are supportive. OSA interventions affected patients and their partners, and decisions regarding treatment may be made within the context of their relationship, with a consideration of the impact of treatment on the spouse.

Although there are no ethical concerns inherent with the technologies used to treat OSA, it can be a challenge for some patients to access some of the necessary resources, such as sleep specialists and specialized sleep labs.

## Considerations

Reimbursement coverage for PAP treatments and oral devices differs across Canada, while surgery may be covered as a medical act. These variations in coverage can be a barrier to accessing effective treatments. Choosing an intervention for specific patients may need to consider the reimbursement criteria, OSA severity, and patient perspectives.

In the economic evaluation, the findings were relatively insensitive to the different reimbursement strategies explored. When oral appliances were expensed out of pocket, they were found to be the most likely cost-effective

intervention for mild-to-moderate OSA. However, a review of patient perspectives and experience and implementation has highlighted the financial burden from out-of-pocket costs as an important issue affecting adherence to treatment.

Trial periods for CPAP with the reuse of devices found to be unacceptable by others may help determine which patients would benefit the most from intervention, without an initial investment. Oral appliances and lifestyle interventions are feasible options for patients for whom CPAP is unacceptable, especially in less severe cases of OSA.<sup>31-33</sup> These alternatives may be appropriate for patients who do not have access to the necessary infrastructure (i.e., electricity, clean water, etc.) for specific treatments.

There was some evidence that the longer the study duration, the lower the effects of CPAP, MADs, and positional therapy, potentially due to discontinuation over time. It is also possible that the effects of CPAP and OAs first peak and then taper, which could also lead to discontinuation over time. In other words, the level of effect first rises and then falls. The authors of one study (retrieved after analysis) concluded that improved clinician communication skills can help support shared decision making and “motivate patients to try CPAP after the initial visit, and thereafter to improve long-term adherence”.<sup>34</sup>

Surgical MMA is invasive and should only be considered if other treatment options have failed, are unacceptable, or are not affordable, and surgery is covered as a medical act. In practice, the appropriate surgical procedure for OSA depends on the site of the anatomical obstruction and a patient’s anatomical features.

Many patients are nonadherent to therapy primarily for personal and contextual reasons. Patients experienced discomfort for all interventions, and this discomfort may change over time as they adjust to the device or recover from surgery. Patients require support from their health care providers and their partners and family. Receiving the right information about treatment choices or how to care for the devices they chose to use is an important component of supporting patients with OSA and their caregivers. Patients felt that it was important to interact with a health care professional following initiation of CPAP, and also expressed a desire for access to professional support and reassurance at night. Further, patients had to persevere with treatment, and the intervention had to become part of their routine for those who could tolerate it.

Although the diagnosis of OSA was beyond the scope of this report, diagnosis is required to access treatment, making access to publicly funded diagnostic testing an important consideration in the treatment of OSA.<sup>35-37</sup> A 2016 study reported that the AHI determined by a device for home diagnosis of sleep apnea was comparable with the results of standard polysomnography. This study suggests that patients may be able to reliably

diagnose sleep apnea at home, possibly increasing access to OSA diagnosis and subsequent treatment.<sup>38</sup>

In addition to differences in clinical presentation, such as sleepiness, fatigue, headache, or mood, men are more commonly reported to be diagnosed with OSA than women.<sup>1</sup> The patient perspectives and experiences review found that one reason for this could be that women are more likely than men to feel shame related to snoring and therefore less likely to seek diagnosis.<sup>39</sup> As well, women may not present with “classic” OSA symptoms,<sup>40</sup> and OSA symptoms do not always correlate with severity. Finally, women are more likely than men to encourage their spouses to be diagnosed, as opposed to the other way around.<sup>41</sup> These findings suggest that OSA could affect women more often than proposed by current diagnosis rates.

## Background

Therapy selection for OSA is based on an assessment of the patient by lab-based polysomnography or home-based portable monitors.<sup>36</sup> An analysis of the clinical effectiveness and cost-effectiveness and a review of patient perspectives and experiences, ethical and implementation issues, and environmental factors were conducted to inform recommendations about the appropriate use of interventions for the treatment of OSA in adults.

The evidence on clinical and economic effectiveness, patient perspectives and experiences, ethical and implementation issues, and environmental factors used for developing this guidance was derived from the CADTH HTA: *Interventions for the Treatment of Obstructive Sleep Apnea in Adults*.<sup>20</sup>

## Research questions

1. What are the clinical effectiveness, comparative clinical effectiveness, and safety concerns of PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle modifications for the treatment of OSA in adults?
  - 1a. What are the clinical effectiveness, comparative clinical effectiveness, and safety concerns of PAP devices, expiratory PAP valve, oral appliances, surgical interventions, and lifestyle for the treatment of adult patients with different OSA severity (i.e., mild, moderate, severe)?
  - 1b. What are the clinical effectiveness, comparative clinical effectiveness, and safety concerns of interventions for the treatment of adult OSA patients with or without comorbidities (e.g., obesity, hypertension, diabetes)?
2. What is the cost-effectiveness of PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle for the treatment of OSA in adults?

3. What are the experiences and perspectives of adult patients, their family members, and their caregivers regarding PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle for the treatment of OSA?
4. What ethical issues are raised by providing PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle to treat OSA in adults? How should these issues be addressed?
5. What are some of the implementation issues associated with PAP devices, expiratory PAP valve, oral appliances, surgical interventions, and lifestyle for the treatment of OSA in adults?
6. What are some potential environmental impacts associated with PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle for the treatment of OSA in adults?

## Summary of The Evidence

### Clinical Evidence

A systematic review of the literature was conducted, using MEDLINE, Embase, Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, Cochrane Central Register of Controlled Trials, and PubMed, for an overview of systematic reviews, meta-analyses, and HTAs, supplemented by a review of primary studies for areas of gap. In total, 33 systematic reviews and 41 primary studies were included in the overview and review, respectively. Both involved adults with OSA who were treated with PAP devices, expiratory PAP valves, oral appliances, surgery, and lifestyle interventions and assessed on various outcomes, with EDS as the primary outcome.

CPAP, expiratory PAP, MADs, TRDs, MMA, GTA, weight loss programs, and positional therapy were all effective at reducing EDS (commonly measured by the Epworth sleepiness scale) compared with inactive controls or pre-treatment. Effect sizes were similar across the interventions, except for people with severe cases of OSA who may benefit more from CPAP than from MADs, although the difference may not be clinically significant. Based on an analysis using OSA severity as the outcome, commonly measured by AHI, effect sizes varied across the interventions, with CPAP showing the largest effect. For people with severe cases of OSA who are eligible for surgery, MMA with or without GTA may be effective at improving both EDS and OSA severity. However, the findings are mostly from small, uncontrolled pre- and post-treatment studies on highly selected patients and warrant caution, considering the invasiveness of the procedure and potential adverse events. Limited evidence was found on other outcomes, such as blood pressure, cardiovascular events, quality of life, and mortality. The 33 systematic reviews and 41 primary studies were assessed to generally be of high quality, using accepted quality assessment tools. But concerns were identified around the study eligibility criteria for the systematic reviews and small samples and uncontrolled pre- and post-treatment study designs for

the primary studies. The primary studies included in the 33 reviews ranged widely in their quality.

## Economic Evidence

A Markov cohort model was constructed to evaluate the cost-effectiveness of various treatment strategies in adult patients diagnosed with OSA (i.e., 76.5% males, 55 years of age) over a patient's lifetime within a Canadian health care payer perspective. The effect of treatment in terms of change in AHI and blood pressure was determined from the clinical review and was translated to changes in the risk of cardiovascular events and motor vehicle accidents in the economic model. The primary outcome was cost per QALY gained in 2016 Canadian dollars. The base-case analysis compared a "no treatment" strategy against PAP therapy, MADs only, and surgery (i.e., MMA with or without GTA). A separate scenario analysis was conducted on obese or overweight patients, in which weight loss would be a suitable treatment strategy.

Cost-effectiveness of treatment strategies for OSA was found to be dependent on a patient's baseline disease severity, as measured by AHI (i.e., lower AHI equates to less severe OSA). At a willingness-to-pay threshold of \$50,000 per QALY, the order in which interventions were considered cost-effective by increasing baseline disease severity was: no treatment (AHI < 15), MADs (15 ≤ AHI ≤ 25), MMA with or without GTA (25 < AHI < 30), PAP therapy (30 ≤ AHI ≤ 32), and MMA with or without GTA (AHI > 32). Absolute gains in QALYs were found to follow a unimodal distribution and were a function of disease severity. Those with mild or more severe OSA had lower gains in QALYs, whereas the largest gains were observed in patients whose baseline severity reduced from severe (AHI ≥ 30) or moderate (15 < AHI < 30) to mild-to-moderate OSA (AHI < 30) or mild OSA (AHI < 15), respectively, due to the impact on subsequent morbidity and mortality risks. Incremental costs were largely driven by the costs of treatment and long-term maintenance costs, given the longer life expectancies of patients on treatment. The model was found to be most sensitive to changes in treatment adherence.

## Patient Perspectives and Experiences Evidence

A systematic review and thematic synthesis of the literature relevant to the research question on patient experience and perspectives were conducted. Patient experience information was identified by searching the following databases: MEDLINE (1946–), Embase (1974–), and PsycINFO (1967–) via Ovid; CINAHL (1981–) via EBSCO; and PubMed. Studies were eligible if they presented the patient or non-clinical caregiver experience. Qualitative studies, surveys, studies with mixed methodology, or systematic reviews of descriptive studies were eligible. A maximum variation approach was used to identify articles for inclusion in the thematic synthesis from a list of eligible articles. A thematic synthesis was conducted, comprising three stages: coding, developing descriptive themes, and developing analytic themes.

Thirty-two studies were included in the thematic synthesis, the coding and analysis of which led to two analytic themes. The first theme described a range of characteristics and factors that influence whether people seek and initiate OSA treatment. Patients are influenced by the information they have on therapy, any disability they may have, whether they receive support for complying with the intervention, and their current life situation. The second analytic theme centred on the finding that interventions for OSA require adaptation to daily routines and relationships; some patients can integrate these interventions into their life and experience benefits, while others are unable to do so. Some patients are noncompliant to therapy for a variety of reasons, each of which is personal and contextual to the individual. Patients reported some degree of discomfort for all interventions, and this discomfort may change over time as patients become accustomed to the device, or recover from surgery.

## Ethical Issues

A systematic review of the normative bioethics literature was conducted to identify literature relevant to the identification and analysis of the potential ethical issues on interventions for OSA (i.e., articles that explicitly and specifically raise ethical issues). Targeted literature searches were performed by a CADTH Information Specialist in MEDLINE, PubMed, and CINAHL from database inception to March 2016. Key terms for ethics concepts and related terms were used and combined with search terms for OSA. The search was limited to English- or French-language literature.

The literature search yielded 1,268 unique citations, none of which passed the first stage of screening because no articles on OSA treatment were found that explicitly mentioned ethical issues. However, in the second stage, the reviewers selected 142 potentially relevant articles that raised implicit ethical issues. Ethical issues relating to OSA were explored according to six key values that emerged from the literature review. The six key values were: respect individual autonomy, maximize benefits and minimize harm for patients, maximize benefits and minimize harms for others affected by OSA, maximize benefits and minimize harms for populations, distribute benefits and burdens of health care resources fairly, and steward scarce resources. In terms of whether universal treatments for OSA should be implemented, they have been shown to offer benefits to OSA patients and reduce overall costs, and so appear to live up to values of conferring a benefit at a population level and stewarding scarce resources. Further, optimizing interventions for OSA to minimize harmful outcomes on both an individual and at the population level is of great benefit, given variability in adherence based on patient behaviours and attitude. To maximize overall benefit, OSA treatment should be provided through an ongoing partnership between health care provider and patient, rather than through discrete events of diagnosis, decision, and intervention.

## Implementation Issues

A narrative literature review was conducted to identify some of the implementation issues associated with the different interventions for the treatment of OSA in adults. Citations arising from the literature searches conducted to address the clinical and economic effectiveness, patient perspectives and experience, and ethical issues were screened independently and in duplicate for information related to implementation issues. Issues identified from relevant studies are organized by OSA intervention (i.e., PAP devices, expiratory PAP valves, oral appliances, surgical interventions, and lifestyle modifications) and further categorized by the level where the issue arises: individual, team, organization, or system or policy. This information was summarized narratively.

From the 27 included studies, one of the biggest implementation issues identified for OSA treatment is difficulty accessing sleep specialists and specialized sleep labs. Home-based portable diagnostic devices and treatment titration options with telehealth-based support are suggested solutions. Most of the implementation evidence focuses on CPAP devices. Barriers to CPAP use include cost and lack of funding as well as patient discomfort or use problems. Suggested CPAP supports include patient education and training, as well as providers and centres that are accredited for the treatment of OSA. Barriers to treatment with oral appliances include lack of physician knowledge and awareness, anatomical and dental health requirements, and the need for regular re-evaluations. Little evidence on implementation issues for OSA surgery or lifestyle interventions was found.

## Environmental Impact

Citations arising from the clinical literature search were screened for information relating to environmental considerations associated with obstructive sleep apnea.

One narrative review article identified the environmental implications associated with OSA. The article briefly examined the environmental considerations of the CPAP unit, including manufacturers adopting green shipping and production methods, creating more energy-efficient products, and using more recyclable materials.

## Research Gaps

Additional research is needed to address patient characteristics that guide the selection of interventions. There is a dearth of evidence on comorbidities and outcomes of the interventions based on patient characteristics. Research on direct, head-to-head comparisons or network meta-analyses on the clinical effectiveness and safety of various treatments, and the impact of these interventions in subgroups of OSA patients who have hypertension or cardiovascular disease on the primary outcome and OSA severity, is also warranted.

Research on OSA treatment for several subgroups, including Indigenous populations, and populations with specific work occupations (e.g., military or law enforcers), is underrepresented in the literature. Further research on these subgroups may provide insight on the most effective treatments, given their living conditions and life situations. Studies on the diagnosis of OSA in women are necessary to assess whether OSA is underdiagnosed, less common, or misdiagnosed among the female population. For instance, there are differences in how men and women describe their symptoms, which may lead to misdiagnosis in women.

Education of primary care clinicians on all the available interventions is necessary, and research on adherence, especially its change with time and relationship with the effectiveness of various treatment interventions, comparative data across treatment interventions, and factors that influence it, is merited. There is also a need for work to evaluate shared decision-making and decision aids in OSA.

## References

1. What is the impact of sleep apnea on Canadians? Fast facts from the 2009 Canadian community health survey - sleep apnea rapid response [Internet]. Ottawa: Public Health Agency of Canada; 2010. [cited 2016 Nov 30]. Available from: <http://www.phac-aspc.gc.ca/cd-mc/sleepapnea-apneesommeil/pdf/sleep-apnea.pdf>
2. Potvin E, Lance JM. Politiques publiques de remboursement d'appareils à pression positive continue pour le traitement de l'apnée obstructive du sommeil [Internet]. Québec (QC): Institut national d'excellence en santé, et en services sociaux; 2014 Apr. [cited 2016 Nov 30]. Available from: [https://www.inesss.qc.ca/fileadmin/doc/INESSS/Rapports/MaladiesRespiratoires/INESSS\\_Politiques\\_remboursement\\_AOS\\_.pdf](https://www.inesss.qc.ca/fileadmin/doc/INESSS/Rapports/MaladiesRespiratoires/INESSS_Politiques_remboursement_AOS_.pdf) . French.
3. Young T, Palta M, Dempsey J, Peppard PE, Nieto FJ, Hla KM. Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study. *WMJ* [Internet]. 2009 Aug [cited 2016 Nov 7];108(5):246-9. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2858234>
4. Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* [Internet]. 2013 May 1 [cited 2016 Nov 7];177(9):1006-14. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3639722>
5. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet*. 2005 Mar 19;365(9464):1046-53.
6. Balk EM, Moorthy D, Obadan NO, Patel K, Ip S, Chung M, et al. Diagnosis and treatment of obstructive sleep apnea in adults [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2011 Jul. [cited 2016 Mar 4]. (AHRQ comparative effectiveness reviews). Available from: [http://www.ncbi.nlm.nih.gov/books/NBK63560/pdf/Bookshelf\\_NBK63560.pdf](http://www.ncbi.nlm.nih.gov/books/NBK63560/pdf/Bookshelf_NBK63560.pdf)
7. Continuous positive airway pressure compared with oral devices or lifestyle changes for the treatment of obstructive sleep apnea: a review of the clinical and cost-effectiveness [Internet]. Ottawa: CADTH; 2014 Dec 22. [cited 2016 Nov 30]. (Rapid response report: summary with critical appraisal). Available from: <https://www.cadth.ca/media/pdf/htis/jan-2015/RC0619%20CPAP%20Final.pdf>
8. Kapur VK. Obstructive sleep apnea: diagnosis, epidemiology, and economics. *Respir Care*. 2010 Sep;55(9):1155-67.
9. CPAP treatment for adults with obstructive sleep apnea: review of the clinical and cost-effectiveness and guidelines [Internet]. Ottawa: CADTH; 2013 Nov 18. [cited 2016 Mar 4]. (Rapid response report: summary with critical appraisal). Available from: <https://www.cadth.ca/sites/default/files/pdf/htis/dec-2013/RC0502%20CPAP%20Treatment%20Final.pdf>
10. Fava C, Dorigoni S, Dalle VF, Danese E, Montagnana M, Guidi GC, et al. Effect of CPAP on blood pressure in patients with OSA/hypopnea: a systematic review and meta-analysis. *Chest*. 2014 Apr;145(4):762-71.
11. National Institute for Health and Care Excellence. Continuous positive airway pressure for the treatment of obstructive sleep apnoea/hypopnoea syndrome [Internet]. London: NICE; 2008 Mar. [cited 2016 Nov 30]. (NICE technology appraisal guidance 139). Available from: <https://www.nice.org.uk/guidance/ta139>
12. Qaseem A, Holty JE, Owens DK, Dallas P, Starkey M, Shekelle P, et al. Management of obstructive sleep apnea in adults: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* [Internet]. 2013 Oct 1 [cited 2016 Mar 4];159(7):471-83. Available from: <http://annals.org/article.aspx?articleid=1742606>
13. Bouloukaki I, Giannadaki K, Mermigkis C, Tzanakis N, Mauroudi E, Moniaki V, et al. Intensive versus standard follow-up to improve continuous positive airway pressure compliance. *Eur Respir J*. 2014 Nov;44(5):1262-74.
14. Wozniak DR, Lasserson TJ, Smith I. Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev*. 2014;1:CD007736.
15. Weaver TE, Grunstein RR. Adherence to continuous positive airway pressure therapy: the challenge to effective treatment. *Proc Am Thorac Soc* [Internet]. 2008 Feb 15 [cited 2016 Oct 6];5(2):173-8. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2645251>
16. Sharples L, Glover M, Clutterbuck-James A, Bennett M, Jordan J, Chadwick R, et al. Clinical effectiveness and cost-effectiveness results from the randomised controlled Trial of Oral Mandibular Advancement Devices for Obstructive sleep apnoea-hypopnoea (TOMADO) and long-term economic analysis of oral devices and continuous positive airway pressure. *Health Technol Assess* [Internet]. 2014 Oct [cited 2016 Mar 4];18(67):1-296. Available from: [http://www.ncbi.nlm.nih.gov/books/NBK262704/pdf/Bookshelf\\_NBK262704.pdf](http://www.ncbi.nlm.nih.gov/books/NBK262704/pdf/Bookshelf_NBK262704.pdf)
17. Dieltjens M, Vanderveken OM, Van den Bosch D, Wouters K, Denollet J, Verbraecken JA, et al. Impact of type D personality on adherence to oral appliance therapy for sleep-disordered breathing. *Sleep Breath*. 2013 Sep;17(3):985-91.
18. Thomasouli MA, Brady EM, Davies MJ, Hall AP, Khunti K, Morris DH, et al. The impact of diet and lifestyle management strategies for obstructive sleep apnoea in adults: a systematic review and meta-analysis of randomised controlled trials. *Sleep Breath*. 2013 Sep;17(3):925-35.
19. Tarasiuk A, Reuveni H. The economic impact of obstructive sleep apnea. *Curr Opin Pulm Med*. 2013 Nov;19(6):639-44.

20. Interventions for the treatment of obstructive sleep apnea in adults: a health technology assessment [Internet]. Ottawa: CADTH; 2017. [cited 2017 Jan 19]. (CADTH optimal use report; vol. 6, no. 1b). Available from: <https://www.cadth.ca/interventions-obstructive-sleep-apnea>
21. Fleetham J, Ayas N, Bradley D, Fitzpatrick M, Oliver TK, Morrison D, et al. Canadian Thoracic Society 2011 guideline update: diagnosis and treatment of sleep disordered breathing. *Can Respir J* [Internet]. 2011 Jan [cited 2016 Mar 4];18(1):25-47. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3070752/>
22. Epstein LJ, Kristo D, Strollo PJ Jr, Friedman N, Malhotra A, Patil SP, et al. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med* [Internet]. 2009 Jun 15 [cited 2016 Mar 18];5(3):263-76. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2699173>
23. Wu H, Yuan X, Zhan X, Li L, Wei Y. A review of EPAP nasal device therapy for obstructive sleep apnea syndrome. *Sleep Breath*. 2015;19(3):769-74.
24. AIM Specialty Health. Sleep disorder management: diagnostic & treatment guidelines [Internet]. Chicago: AIM Specialty Health; 2015 Aug 27. [cited 2016 Mar 18]. Available from: <http://www.aimspecialtyhealth.com/clinical-guidelines/agreementAccess-sleep>
25. Obstructive sleep apnea: the role of dentists in the treatment of snoring and obstructive sleep apnea with oral appliances [Internet]. Vancouver: College of Dental Surgeons of British Columbia; 2014 Jun. [cited 2016 Mar 18]. (CDSBC standards and guidelines). Available from: <https://www.cdsbc.org/CDSBCPublicLibrary/Obstructive-Sleep-Apnea.pdf>
26. Boyd SB, Walters AS, Song Y, Wang L. Comparative effectiveness of maxillomandibular advancement and uvulopalatopharyngoplasty for the treatment of moderate to severe obstructive sleep apnea. *J Oral Maxillofac Surg* [Internet]. 2013 Apr [cited 2016 May 27];71(4):743-51. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3604163>
27. Caples SM, Rowley JA, Prinsell JR, Pallanch JF, Elamin MB, Katz SG, et al. Surgical modifications of the upper airway for obstructive sleep apnea in adults: a systematic review and meta-analysis. *Sleep* [Internet]. 2010 Oct [cited 2016 Nov 30];33(10):1396-407. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2941427>
28. Song SA, Chang ET, Certal V, Del Do M, Zaghi S, Liu SY, et al. Genial tubercle advancement and genioplasty for obstructive sleep apnea: a systematic review and meta-analysis. *Laryngoscope*. 2016 Aug 22. Epub ahead of print.
29. Emara TA, Omara TA, Shouman WM. Modified genioglossus advancement and uvulopalatopharyngoplasty in patients with obstructive sleep apnea. *Otolaryngol Head Neck Surg*. 2011 Nov;145(5):865-71.
30. Kim SY, Park JE, Lee YJ, Seo HJ, Sheen SS, Hahn S, et al. Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *J Clin Epidemiol*. 2013 Apr;66(4):408-14.
31. Iftikhar IH, Bittencourt L, Youngstedt SD, Ayas N, Cistulli PA, Schwab R, et al. Comparative efficacy of CPAP, MADs, exercise-training and dietary weight loss for sleep apnea: a network meta-analysis. *Sleep Med*. 2016 Jun 28. Epub ahead of print.
32. Bratton DJ, Gaisl T, Schlatzer C, Kohler M. Comparison of the effects of continuous positive airway pressure and mandibular advancement devices on sleepiness in patients with obstructive sleep apnoea: a network meta-analysis. *Lancet Respir Med*. 2015 Nov;3(11):869-78.
33. Bratton DJ, Gaisl T, Wons AM, Kohler M. CPAP vs mandibular advancement devices and blood pressure in patients with obstructive sleep apnea: a systematic review and meta-analysis. *JAMA*. 2015 Dec 1;314(21):2280-93.
34. Brostrom A, Fridlund B, Hedberg B, Nilsson P, Ulander M. Communication between patients with obstructive sleep apnea syndrome and health care personnel during the initial visit to a CPAP clinic. *J Clin Nurs*. 2016 Sep 29. Epub ahead of print.
35. Alshaer H, Fernie GR, Tseng WH, Bradley TD. Comparison of in-laboratory and home diagnosis of sleep apnea using a cordless portable acoustic device. *Sleep Med*. 2016 Jun;22:91-6.
36. Health Technology and Policy Unit, School of Public Health, Department of Public Health Sciences, University of Alberta. Level I and level III sleep studies for the diagnosis of sleep disordered breathing (SDB) in adults: final report [Internet]. Edmonton: Alberta Health; 2013 Jul 25. [cited 2016 Nov 30]. Available from: <http://www.health.alberta.ca/documents/AHTDP-Sleep-UofA-2013.pdf>
37. El Shayeb M, Topfer LA, Stafinski T, Pawluk L, Menon D. Diagnostic accuracy of level 3 portable sleep tests versus level 1 polysomnography for sleep-disordered breathing: a systematic review and meta-analysis. *CMAJ* [Internet]. 2014 Jan 7 [cited 2016 Nov 21];186(1):E25-E51. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883848>
38. Aurora RN, Casey KR, Kristo D, Auerbach S, Bista SR, Chowdhuri S, et al. Practice parameters for the surgical modifications of the upper airway for obstructive sleep apnea in adults. *Sleep* [Internet]. 2010 Oct [cited 2016 Mar 18];33(10):1408-13. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2941428>
39. Broström A, Johansson P, Strömberg A, Albers J, Martensson J, Svanborg E. Obstructive sleep apnoea syndrome--patients' perceptions of their sleep and its effects on their life situation. *J Adv Nurs*. 2007 Feb;57(3):318-27.
40. Lin CM, Davidson TM, Ancoli-Israel S. Gender differences in obstructive sleep apnea and treatment implications. *Sleep Med Rev*. 2008 Dec;12(6):481-96.
41. Henry D, Rosenthal L. "Listening for his breath:" the significance of gender and partner reporting on the diagnosis, management, and treatment of obstructive sleep apnea. *Soc Sci Med*. 2013 Feb;79:48-56.

## Appendix 1: HTERP

HTERP consists of up to seven core members appointed to serve for all topics under consideration during their term of office, and up to five expert members appointed to provide their expertise for a specific topic. For this project, three expert members were appointed; their expertise included internal medicine, clinical chemistry, pathology, and family medicine. The core members include health care practitioners and other individuals with expertise and experience in evidence-based medicine, critical appraisal, health technology assessment, bioethics, and health economics. One public member is also appointed to the core panel to represent the broad public interest.

HTERP is an advisory body to CADTH and is convened to develop guidance or recommendations on non-drug health technologies to inform a range of stakeholders within the Canadian health care system. Further information regarding HTERP is available at [www.cadth.ca/en/advisory-bodies/health-technology-expert-review-panel](http://www.cadth.ca/en/advisory-bodies/health-technology-expert-review-panel).

### HTERP Core Members

Dr. Stirling Bryan (Chair)

Dr. Jenny Basran

Dr. Leslie Anne Campbell

Dr. Hilary Jaeger

Dr. Jeremy Petch

Dr. Lisa Schwartz

Ms. Tonya Somerton

### Expert Members

Dr. Sachin Pendharkar

Dr. Najib Ayas

Dr. Jessica Otte

### Conflict of Interest

No members declared any conflicts of interest. [Conflict of Interest Guidelines](#) are posted on the CADTH website.

## Appendix 2: HTERP Deliberative Framework

**Table 1: HTERP Deliberative Framework for Interventions for Obstructive Sleep Apnea**

Framework Domain	Examples of Information and Element(s)	Possible HTERP Discussion Question(s)	Discussion Points
<b>Background/Context</b>	Audience; issue and policy question(s)	Who requested this assessment?  Why?	<ul style="list-style-type: none"> <li>• Need for evidence-informed selection of OSA intervention based on patient characteristics.</li> <li>• Identify considerations for prioritization of patients, when device supplies are limited.</li> <li>• Implementation of an efficient, respectful, and equitable OSA care pathway.</li> </ul>
<b>Need</b>	Background on health condition  Size of affected population	What condition does this health technology address?  How many patients could potentially be affected?	<ul style="list-style-type: none"> <li>• 3% of adult population is diagnosed with OSA; 17% to 25% are suspected to be or are at risk (i.e., age, male sex, obesity risk factors).</li> </ul>
	Availability of alternatives	Are there existing therapeutic/diagnostic technologies that address the same problem?	<ul style="list-style-type: none"> <li>• PAP reimbursement varies across jurisdictions (i.e., partial to complete coverage).</li> <li>• Dental devices coverage varies.</li> <li>• Surgery may be covered as a medical act.</li> </ul>
<b>Benefits</b>	Efficacy  Clinical effectiveness  Impact on patient-centred outcomes  Impact on clinical management  Non-health benefits (e.g., patient autonomy, dignity)	Has the clinical effectiveness of the candidate technology been established?  Compared with what?  What improvements does this technology purport to offer over others?  What types of evidence is this based on?  Are we aware of any better-quality evidence likely to be produced in the near future?	<ul style="list-style-type: none"> <li>• For severe OSA, CPAP more effective than MADs for excessive daytime sleepiness.</li> <li>• For severe OSA, MMA ± GTA most effective if PAP therapies failed (Note: outcomes were derived from small, uncontrolled studies).</li> <li>• For other OSA severities, all interventions similarly effective.</li> <li>• CPAC is superior for reducing OSA severity.</li> <li>• Limited evidence on blood pressure, cardiovascular events, QoL, mortality outcomes is available.</li> </ul>

Framework Domain	Examples of Information and Element(s)	Possible HTERP Discussion Question(s)	Discussion Points
		Are there any non-health benefits?	<ul style="list-style-type: none"> <li>Some literature is of high quality but there are concerns with regards to the eligibility criteria, small samples, and uncontrolled studies.</li> </ul>
<b>Harms</b>	Safety	<p>What is known about safety in absolute terms, and in comparison with existing technologies?</p> <p>What types of evidence is this based on?</p>	<ul style="list-style-type: none"> <li>No major adverse events were reported, except MMA + GTA, with which all patients had complications caused by infections.</li> </ul>
<b>Patient Preferences</b>	Acceptability of health technology by the patient	<p>How will it potentially affect patients and what are their opinions about the technology?</p> <p>How acceptable is it to patients?</p>	<ul style="list-style-type: none"> <li>Patient factors are individualized and contextual.</li> <li>OSA affects bed partners. Patients perform a trade-off between the benefits of using interventions and their discomfort.</li> <li>Elements of support and information are needed.</li> <li>No evidence on patient experience with surgery was identified.</li> </ul>
<b>Economic Impact</b>	<p>Cost-effectiveness</p> <p>Infrastructure support costs</p> <p>Budget impact</p>	<p>What will the technology cost (including initial purchase price and consumables, maintenance, and training of personnel)?</p> <p>Is there evidence of value for money? How is value defined?</p> <p>What is the expected lifespan and total budget impact of the technology?</p>	<ul style="list-style-type: none"> <li>PAP or MMA is the most cost-effective for severe OSA with a baseline AHI at a WTP of \$50,000/QALY.</li> <li>MAD or MMA is most cost-effective for moderate OSA.</li> <li>No treatment is most cost-effective for mild OSA.</li> </ul>
<b>Implementation</b>	<p>Integration of technology into existing workflow</p> <p>Training/competency requirements</p> <p>Repair and maintenance</p>	<p>Have issues of implementation of the technology in a real-world health system been identified and addressed?</p>	<ul style="list-style-type: none"> <li>One of the biggest implementation issues for OSA treatment is the difficulties in accessing sleep specialists and specialized sleep labs.</li> <li>Most of the implementation evidence focuses on CPAP.</li> <li>CPAP barriers include cost or lack of funding, patient discomfort or difficulties using the device.</li> </ul>

Framework Domain	Examples of Information and Element(s)	Possible HTERP Discussion Question(s)	Discussion Points
			<ul style="list-style-type: none"> <li>• CPAP enablers encompass patient education, training, or support accredited sleep centres or health care professionals, but there are no accepted guidelines.</li> <li>• OA barriers include lack of knowledge or awareness, anatomical and dental health requirements, need for regular re-evaluations.</li> <li>• OA enablers encompass multidisciplinary sleep clinics.</li> <li>• Little evidence on implementation issues for OSA surgery or lifestyle interventions was found.</li> </ul>
<b>Legal</b>	Legal impacts	Are there potential legal or regulatory aspects to the introduction and use of this technology?	<ul style="list-style-type: none"> <li>• The legal issues associated with driving were discussed. For example, some jurisdictions can revoke a driver's licence if the person has severe OSA.</li> <li>• The Canada Health Act does not include dentists.</li> </ul>
<b>Ethics</b>	Consistent with Canadian ethical values	<p>Are there potential issues of equity (access by particular populations, for example) with respect to introducing this technology?</p> <p>Are there any other ethical issues to consider?</p>	<ul style="list-style-type: none"> <li>• Interventions for OSA do not appear to present ethical concerns that are inherent to the technologies.</li> <li>• Several values are relevant on how access to OSA interventions are organized and delivered.</li> <li>• Duties to act in ways that maximize benefits to patients or others, respect patient choice, and ensure reasonable access to resources are of core importance.</li> <li>• The capacity for patients to benefit from most OSA interventions relies heavily on the patient's behaviour, so patient context is perhaps unusually significant for these technologies.</li> </ul>
<b>Environmental Impact</b>	Environmental impact of health technology	What is the potential impact on the environment of this technology?	<ul style="list-style-type: none"> <li>• One review article examined the environmental considerations of the CPAP unit including:               <ul style="list-style-type: none"> <li>○ Green shipping and production methods</li> </ul> </li> </ul>

Framework Domain	Examples of Information and Element(s)	Possible HTERP Discussion Question(s)	Discussion Points
			<ul style="list-style-type: none"> <li>○ Creating more energy-efficient products</li> <li>○ Using more recyclable materials.</li> </ul>
<b>Other</b>		<p>Are there particular questions with regards to professional fees that have been identified and addressed?</p> <p>Does this candidate technology raise some questions that are not addressed by the above set of questions?</p>	<ul style="list-style-type: none"> <li>● Compliance, adherence, and patient acceptability of the various treatment options.</li> <li>● Accurate diagnosis is necessary before appropriate treatment can be given.</li> </ul>

AHI = apnea-hypopnea index; CPAP = continuous positive airway pressure; GTA = genial tubercle advancement; HTERP = Health Technology Expert Review Panel; MAD = mandibular advancement device; MMA = maxillomandibular advancement; OA = oral appliance; OSA = obstructive sleep apnea; PAP = positive airway pressure; QALY = quality-adjusted life year; QoL = quality of life; WTP = willingness to pay.